

WETLAND INVENTORY

KANGAROO ISLAND

An assessment of selected inland wetlands
of Kangaroo Island, South Australia.



Government
of South Australia

This report may be cited as:

Seaman, R.L. (2002) *Wetland Inventory of Kangaroo Island*.
South Australia. Department for Environment and Heritage.

Author

Russell Seaman
Scientific Officer, Wetlands Management.
National Parks and Wildlife

Cartography and design

Russell Seaman

Photographs

Russell Seaman

Geographical Information System Data

Supplied by Geographical Analysis and Research Unit, Planning SA,
Department for Transport, Urban Planning and the Arts.

and

The Department for Environment and Heritage, South Australia.

Funding Sources

Environment Australia

National Parks and Wildlife SA

Acknowledgments

In the preparation of this report the expertise, advice and support of several people is greatly appreciated.

- Rob Walsh
- Roman Urban, DEH.
- Sandy Carruthers, Planning SA.
- Tim Croft, DEH.
- Kylie Moritz, DEH.

Executive Summary

During November and December 2000 a wetland inventory was undertaken on Kangaroo Island which documented a representative sample of inland wetlands by recording their physical, chemical and biological attributes. Twenty-six wetlands were surveyed and included saline lake systems (19 sites), brackish water bodies and freshwater wetlands. Breakneck River and Six-mile Lagoon are classified as the only freshwater wetlands surveyed.

The aquatic invertebrate fauna was notably scarce in many of the salt lakes surveyed, however, 14 wetlands displayed good invertebrate trophic levels. The correlation between increasing salinity levels and decreasing biological activity was clear; this decline is of concern for the health of many wetlands on Kangaroo Island.

Sixteen wetlands surveyed are considered to be nationally important by meeting the ANZECC criteria of being a good example of a wetland type occurring within a biogeographic region in Australia. These wetlands include Six-mile Lagoon, Duck Lagoon and Destree Swamp. Three wetlands are recommended for monitoring; these are Grassdale lagoon, Six-mile Lagoon and Rush Lagoon.

SECTION ONE - WETLAND INVENTORY.....	9
1.0 INTRODUCTION	9
2.0 REPORT STRUCTURE.....	9
3.0 PROJECT SCOPE	9
4.0 OVERVIEW OF PAST WETLAND INVENTORY STUDIES.....	10
5.0 WETLAND RISK ASSESSMENT	11
6.0 WETLAND INVENTORY METHODOLOGY	12
6.1 STUDY AREA BOUNDARIES	12
6.2 SITE SELECTION.....	12
6.3 GIS DATABASE.....	12
6.4 WETLAND INVENTORY SURVEY.....	13
7.0 WETLAND SURVEY TEMPLATE.....	13
SECTION TWO - WETLAND ASSESSMENT FOR KANGAROO ISLAND.....	14
8.0 INTRODUCTION	14
8.1 WETLAND OVERVIEW.....	14
9.0 WETLAND LAND USE	15
9.1 BACKGROUND.....	15
9.2 ANALYSIS	15
10.0 TENURE AND MANAGEMENT AUTHORITY.....	16
10.1 BACKGROUND.....	16
10.2 ANALYSIS	16
11.0 ENVIRONMENTAL ASSOCIATIONS AND IBRA REGIONS	17
11.1 BACKGROUND.....	17
11.2 ANALYSIS	17
12.0 KEY BIODIVERSITY AREAS.....	18
12.1 BACKGROUND.....	18
12.1 ANALYSIS	19
13.0 WETLAND AREA.....	19
13.1 BACKGROUND.....	19
13.2 ANALYSIS	19
14.0 LANDFORM ELEMENT	20
14.1 BACKGROUND.....	20
14.2 ANALYSIS	20
15.0 GEOLOGY.....	21
15.1 BACKGROUND.....	21
15.1 ANALYSIS	21
16.0 HYDROLOGY.....	22
16.1 BACKGROUND.....	22
16.2 ANALYSIS	22
17.0 FLORA ANALYSIS	23

18.0	DEGRADATION AND DISTURBANCE	26
18.1	BACKGROUND	26
18.2	ANALYSIS	26
19.0	AQUATIC VEGETATION CLASSES	27
19.1	BACKGROUND	27
19.2	ANALYSIS	27
20.0	AQUATIC FAUNA ANALYSIS	29
20.1	INVERTEBRATES	29
20.1.1	<i>Ecological benefits</i>	29
20.1.2	<i>Trophic dynamics</i>	29
20.2	TEMPORARY WETLANDS	30
20.3	SALINE SYSTEMS	31
20.4	ANALYSIS	31
20.4.1	<i>Frequency of invertebrate occurrence</i>	31
20.4.2	<i>Trophic levels</i>	34
20.5	AMPHIBIANS	38
20.6	FISH	38
20.7	MAMMALS	38
20.8	AVI-FAUNA	38
21.0	WATER CHEMISTRY	39
21.1	PH	40
21.1.1	<i>Background</i>	40
21.1.2	<i>Analysis</i>	41
21.2	CONDUCTIVITY	41
21.2.1	<i>Background</i>	41
21.1.2	<i>Analysis</i>	42
21.3	TURBIDITY	44
21.3.1	<i>Background</i>	44
21.3.2	<i>Analysis</i>	46
21.4	WATER TEMPERATURE	46
21.4.1	<i>Background</i>	46
21.4.1	<i>Analysis</i>	47
21.5	DISSOLVED OXYGEN	47
21.5.1	<i>Background</i>	47
21.5.2	<i>Analysis</i>	48
22.0	RAPID ASSESSMENT	49
22.1	AQUATIC FAUNA	49
22.2	AQUATIC FLORA	50
22.3	RIPARIAN VEGETATION RAPID ASSESSMENT	51
22.3.1	<i>Analysis - Toe of bank</i>	51
23.3.2	<i>Bank</i>	52
23.3.3	<i>Buffer</i>	52
23.0	WETLAND CONDITION SCORE	53
24.0	RAPID ASSESSMENT TOTAL SCORE	54
25.0	ANZECC WETLAND CRITERIA	55
25.1	ANALYSIS	55
26.0	WETLAND TYPES	57
26.1	<i>Analysis</i>	58

SECTION THREE - WETLAND MONITORING	59
27.0 INTRODUCTION	59
27.1 <i>Ecological change</i>	59
28.0 MONITORING PROTOCOLS AND INDICATORS	61
28.1 EARLY WARNING INDICATORS.....	62
28.2 TOXICITY TESTS	62
28.3 FIELD EARLY WARNING INDICATORS.....	63
28.4 RAPID ASSESSMENTS.....	63
29.0 RECOMMENDED INDICATORS FOR MONITORING SURVEYED WETLANDS.....	63
29.1 EARLY WARNING INDICATORS – WATER CHEMISTRY.....	63
29.2 ECOSYSTEM BASED INDICATOR	64
29.3 OTHER CONSIDERATIONS	64
30.0 RECOMMENDED WETLANDS TO MONITOR	65
30.1 RECOMMENDED PRIORITY WETLANDS TO MONITOR	65
31.0 RECORDING MONITORING PARAMETERS	66
BIBLIOGRAPHY	67
APPENDIX 1 WETLAND INVENTORY DESCRIPTIONS.....	71
APPENDIX 2 INVERTEBRATE RECORDS FOR SURVEYED WETLANDS.....	82

Figures

FIGURE 9-1. ON SITE WETLAND LAND USES	15
FIGURE 11-1. NUMBER OF WETLANDS LOCATED IN ENVIRONMENTAL ASSOCIATIONS	17
FIGURE 12-1. WETLANDS WITHIN KEY BIODIVERSITY AREAS	18
FIGURE 13-1. WETLAND AREA	19
FIGURE 14-1. WETLAND LANDFORM ELEMENTS	20
FIGURE 19-1. AQUATIC VEGETATION TYPES	27
FIGURE 20-1. INVERTEBRATE SCORES	34
FIGURE 21-1. TEMPORARY WETLAND CYCLES	39
FIGURE 21-2. PH VALUES.....	40
FIGURE 21-3. FISH ACTIVITY AGAINST TURBIDITY VALUES AND TIME	45
FIGURE 22-1. AQUATIC FAUNA RAPID ASSESSMENT	49
FIGURE 22-2. AQUATIC FLORA RAPID ASSESSMENT	50
FIGURE 22-3. RAPID ASSESSMENT, LOW – HIGH WATER.....	51
FIGURE 22-4. RAPID ASSESSMENT, BANK.....	52
FIGURE 22-4. RAPID ASSESSMENT, BUFFER	52
FIGURE 23-1. WETLAND SCORE.....	53
FIGURE 24-1. RAPID ASSESSMENT SCORES	54
FIGURE 26-1. WETLAND TYPES IN KANGAROO ISLAND.....	58

Plates

PLATE 1. DUCK LAGOON.....	15
PLATE 2. SIX MILE LAGOON 2.....	16
PLATE 3. LUNETTE ADJACENT TO WHITE LAGOON.....	21
PLATE 4. SIX MILE LAGOON.....	22
PLATE 5. HALLS ROAD SWAMP	25
PLATE 6. DESTREE SWAMP	25
PLATE 7. CYGNET RIVER LAGOON.....	25
PLATE 8. HALLS ROAD SWAMP	26
PLATE 9. RUSH LAGOON	28
PLATE 10. GRASSDALE LAGOON	28
PLATE 11. DISCOVERY LAGOON.....	28
PLATE 12. DIACYPRIS CF.SPINOSA.....	32
PLATE 13. <i>DICROTENDIPES</i> SP.	32
PLATE 14. <i>TRICOPLEURA OECETIS</i>	32
PLATE 15. CHIRONOMIDAE	36
PLATE 16. ANISOPS.....	36
PLATE 17. SIX MILE LAGOON.....	42
PLATE 18. DUCK LAGOON.....	43
PLATE 19. BIG WHITE SALT LAGOON.....	43
PLATE 20. SALT LAKE.....	43
PLATE 21.. NEPEAN BAY LAGOON	47
PLATE 22. RUSH LAGOON.....	48
PLATE 23. CYGNET RIVER LAGOON.....	48
PLATE 24. LASHMAR LAGOON.....	51
PLATE 25. GRASSDALE LAGOON	53
PLATE 26. BREAKNECK RIVER	56
PLATE 27. GRASSDALE LAGOON.....	56
PLATE 28. BAY LAKE.....	58
PLATE 29. EMANUAL LAGOON.....	58

Maps

- Map 1** **Kangaroo Island wetland study area**
- Map 2** **Surveyed wetlands on Kangaroo Island**
- Map 3** **Environmental Regions on Kangaroo Island**
- Map 4** **Key Biodiversity Areas on Kangaroo Island**
- Map 5** **Geological Features underlying surveyed wetlands**

SECTION ONE WETLAND INVENTORY

1.0 INTRODUCTION

Kangaroo Island is one of the few remaining regions in South Australia with inland temperate freshwater wetlands in near natural condition. The freshwater wetlands on Kangaroo Island are unique because many are independent from river systems forming their own hydrological unit within the landscape.

This Wetland Inventory of Kangaroo Island documents a representative sample of inland wetlands and records their physical, chemical and biological attributes. From this information, aquatic environments that contain high biodiversity are highlighted, and threats documented. The inventory provides a snap shot of the condition and conservation value of wetlands throughout Kangaroo Island. Those wetlands that meet one or more of the Australian and New Zealand Environment and Conservation Council (ANZECC) criteria for an important wetland will be nominated for inclusion into the National Directory of Important Wetlands in Australia.

The Wetland Inventory for Kangaroo Island is an initiative of the Wetland Section-Environment Australia and the South Australian Department for Environment and Heritage.

2.0 REPORT STRUCTURE

This report is divided into three sections, namely of the wetland inventory, wetland assessment and wetland monitoring.

Section 1 - Wetland Inventory. Outlines the project aims and inventory methodology.

Section 2 - Wetland Assessment. Provides an analysis of the wetland inventory, which includes the identification of wetland values and threats.

Section 3 – Wetland Monitoring. Discusses frameworks for monitoring, and recommends indicator species for monitoring and specific wetlands to monitor.

3.0 PROJECT SCOPE

The project scope consist of five actions, these are to:

1. Undertake baseline wetland surveys for inland surface waters in on Kangaroo Island.
2. Identify wetlands of conservation significance, according to agreed ANZECC classification.
3. Provide digital coverage of spatial boundaries of identified wetlands in ARC/INFO compatible format at appropriate scale, in consultation with Planning SA.
4. Produce a report detailing the physical, biological and chemical attributes of wetlands on Kangaroo Island.
5. Nominate wetlands of significance for inclusion in the Directory of Important Wetlands in Australia.

4.0 OVERVIEW OF PAST WETLAND INVENTORY STUDIES

The most comprehensive listing of wetlands in South Australia in terms of numbers and coverage is by Lloyd and Balla (1986). This study identified approximately 1500 wetlands and complexes state-wide.

The Lloyd and Balla listing was a desktop study that collated and recorded information within a standard format. This included:

- wetland type
- name
- location
- size
- catchment
- aquatic fauna
- aquatic and fringing vegetation
- wetland condition
- water regime
- landuse
- impacts
- tenure .

The Lloyd and Balla report provides a good starting point in understanding the extent and some attributes of South Australian wetlands. However, the study falls short in providing up to date information on invertebrate composition, water chemistry and basic landform information. Fifty-three wetlands are documented for Kangaroo Island in the Lloyd and Balla report, this includes swamps, lakes and rivers.

In 1984, Barritt completed a study of the water resources on Kangaroo Island. Barritt describes six different groups of lagoons on Kangaroo Island and surveyed a total of 36 lagoons, 14 river sites and five dryland areas. His report provides a general overview of wetland resources through field notes and observations.

Since Lloyd and Balla's 1986 report, several state-wide studies have mirrored this kind of information collection and presentation, but have generally not collected detailed field baseline wetland information. None-the-less, good information has been generated for certain areas including the Murray River corridor, for example, Thompson's (1986) study documented, mapped and described the River Murray Wetlands and Jensen *et al* (1996) *Wetland Atlas Report of the South Australian Murray Valley Wetlands* made inroads into spatially capturing the locations through the use of GIS. The introduction of linking wetlands with GIS enabled the creation of a wetlands GIS database for the Murray Valley Wetlands. In 1997 Carruthers and Hille developed a GIS database for the South East wetlands. This database recorded wetland type, name, complex, watercourses and assigned a condition score and conservation value. The benefits of collecting data and linking it to GIS became evident not only for environmental planning and for information retrieval, but also for reporting to Environment Australia on the extent of wetland resources.

In 1993 the Australian Nature and Conservation Agency published the first edition of '*A Directory of Important Wetlands in Australia*'. A second edition was compiled in 1996, which included information on 68 wetlands in South Australia, 13 of which are located on Kangaroo Island. This constitutes 72% of the important wetlands in the Mount Lofty Block IBRA region, (Willoughby, N. *et al* 2001)

De Jong and Morelli (1996) commented that in compiling the Directory of Important Wetlands it became apparent that several regions within South Australia were lacking baseline wetland information. They suggested that there is a need for systematic inventories, biological surveys and research programs in many areas of the State. Wetland information in regions such as the Great Victoria Desert, Flinders and Olary Ranges, Eyre Peninsula, Yorke Peninsula, Kangaroo Island and Nullarbor are inadequate.

The Wetland Inventory for the Kangaroo Island combines some of these recommendations, including continuing the development of a GIS database and providing baseline information which includes physical, chemical and biological information for wetlands on Kangaroo Island.

5.0 WETLAND RISK ASSESSMENT

The wetland risk assessment is a conceptual framework to assist in predicting and assessing change in the ecological character of wetlands. The framework has been adopted by Ramsar (resolution V11.10) and is now promoted as an integral component of the management planning process for wetlands. The relevance of undertaking wetland inventories becomes apparent within this framework. A wetland inventory ultimately collects information for the wetland assessment framework. This information is also critical in order to make recommendations for monitoring.

A central component of the wetland risk assessment is the ability to record the ecological character of a wetland. The ecological character is the sum of the biological, physical, and chemical components of the wetland ecosystem and their interactions that maintain wetland functions and attributes. Recording changes to the ecological character of a wetland involves the development of a monitoring program, this is further discussed in Section Three – Wetland Monitoring.

6.0 WETLAND INVENTORY METHODOLOGY

A wetland inventory essentially collects information that assists in wetland management. It can provide information for specific assessment and monitoring activities.

6.1 Study area boundaries

The project boundary is based on the Kangaroo Island Natural Heritage Trust (NHT) boundary. Refer to [Map 1](#).

6.2 Site selection

The aim of the wetland selection process was to sample a broad range of wetlands across Kangaroo Island. Factors such as time constraints, accessibility and the project budget were limiting factors in the number of wetlands selected. Wetlands were selected initially by studying the GIS waterbody coverage for Kangaroo Island. Waterbodies within State government lands, community lands, Council managed lands, land under management agreements and Heritage Agreements were targeted for further investigation. Stakeholders such as National Parks and Wildlife were also consulted.

A total of 26 wetlands were surveyed on Kangaroo Island; refer to [Map 2](#) for wetland locations.

6.3 GIS Database

This project builds on initiatives undertaken by Planning SA and the Department for Environment and Heritage. GIS databases have been developed for the Murray River region by Carruthers and Nicolson (1992) and published in the form of an Atlas, Jensen *et al* (1996). A GIS database exists for the South East region of the state and has been published in the form of a technical report by Carruthers and Hille (1997). No other regions in South Australia have a GIS wetland database at present. One of the project objectives is to provide a digital coverage of spatial boundaries of wetlands surveyed on Kangaroo Island.

A state wide numbering system was developed for identifying wetlands which follows the system established for the Murray River wetlands. The Murray River wetlands have been assigned the numbers S0001 to S0999. The South East region (numbers S1000 to S1999), Eyre Peninsula, (numbers S3000 to S3999), Northern Agricultural Districts numbers S4000 to S4999 and the Mount Lofty Ranges (numbers S2000 to S2999). Kangaroo Island has been assigned the numbers S5000 to S5999. The software used to produce the wetlands data is the ESRI (Environmental Systems Research Institute) geographic information system (GIS) ARC/INFO. The GIS layer was created initially from the existing land cover layer that contained areas designated as swamps, vegetated swamps, lakes and vegetated lakes. This land cover layer was mapped from 1:40 000 colour aerial photography by the Geographical Analysis and Research Unit, Planning SA, Department for Transport, Urban Planning and the Arts.

6.4 Wetland inventory survey

In developing the wetland survey, it was critical that information collected could be used for an initial assessment of wetland character. This ultimately involved the collection of physical, biological and chemical parameters. In the development of the survey template, several methodologies were studied and adapted, these include:

- Butcher, R.J. (1999) *Assessing biodiversity in temporary and permanent wetlands*. pp. 50-53 in *The Other 99%. The Conservation and Biodiversity of Invertebrates*, ed by Ponder, W. and Lunney, D. (1999). Transactions of the Royal Zoological Society of New South Wales.
- Finlayson, C.M. and Spiers, A.G. (1999). *Techniques for enhanced wetland inventory and monitoring*. Supervising Scientist, Canberra
- Fairweather, P.G. and Napier (1998). *Environmental indicators for national state of the environment reporting - inland waters*. Environment Australia.
- Maher, W. and Liston, P. (1997). *Water quality for maintenance of aquatic ecosystems: Appropriate indicators and analysis. Australia: State of the Environment Technical Paper Series. (Inland waters)*. Environment Australia.
- Morelli, J. and de Jong, M. (1996). *A Directory of Important Wetlands in South Australia*. South Australian Department of Environment and Natural Resources, Adelaide.
- Storey, A.W., Lane, J.A.K. and Davies, P.M. (1997). *Monitoring the ecological character of Australia's wetlands of international importance (RAMSAR Convention)*. Western Australian Department of Conservation and Land Management and Biodiversity Group of Environment Australia.

7.0 WETLAND SURVEY TEMPLATE

For each wetland surveyed physical, biological and chemical information was collected, a brief outline is given below. The complete wetland survey descriptions are given in Appendix 1.

Physical parameters

- Wetland Reference Number
- Ramsar Site
- Land use
- Land element
- Wetland name
- Description of site
- Tenure
- Geology

Biological parameters

- Vegetation associations
- Biological threats
- Noteworthy flora and fauna
- Aquatic vegetation classes

Chemical parameters

- Dissolved oxygen
- Conductivity
- Turbidity
- pH
- Temperature

SECTION TWO WETLAND ASSESSMENT FOR KANGAROO ISLAND

8.0 INTRODUCTION

Wetland assessment involves the identification, cataloguing of values and status of threats to wetlands as a basis for the collection of more specific information through monitoring. This section presents each survey parameter by discussing background information and analysis of data collected during field assessment.

8.1 Wetland overview

The Wetland Inventory surveyed two main types of inland wetlands; these are classified as saline wetlands and freshwater wetlands. The saline wetlands generally occur on the eastern section of Kangaroo Island (within the Mac Gillivray and Amberley environmental associations). These saline wetlands are located in a largely fragmented habitat with cleared land occupying a majority of the landscape. A large concentration of saline wetlands occur approximately 20 kilometres south from Kingscote. The largest of these wetlands are Rush Lagoon, White Lagoon and Waidrowski Lagoon.

The freshwater wetlands tend to be located on the western section of Kangaroo Island with wetlands of high biodiversity values located within the National Parks and Wildlife reserve areas or areas under private conservation management. Wetlands such as Grassdale Lagoon, Six Mile lagoon and further west, Murray's Lagoon are excellent examples of inland freshwater wetlands.

[Map 2](#) illustrates the wetland survey locations on Kangaroo Island.

9.0 WETLAND LAND USE

9.1 Background

Land use around and in wetland areas usually dictates the level of protection and the condition of the wetland. This wetland survey is biased toward lands that have a land use favouring conservation, as these tend to contain wetlands with high biodiversity values. Surrounding land uses are a major contributor to the processes that may threaten wetlands. Many wetlands on Kangaroo Island are under threat from increasing resulting from the clearance of native vegetation.

9.2 Analysis




Figure 9-1. On site wetland land uses

Note: these figure include 15 sites, which were not included in the wetland survey, but information was included from Lloyd and Balla (1986) and Barritt (1982).

The importance of off-reserve wetland conservation becomes evident with 15 sites recording a land use of private conservation. Several sites (such as Rush Lagoon) are within private land and under a Heritage Agreement. Eleven wetlands surveyed have a primary land use of grazing (eg Lashmar Lagoon) and eight sites have a conservation management land use being in NPWSA reserves (eg Grassdale Lagoon). Seven sites recorded a land use of cereal and legume production.



Plate 1. Duck lagoon

Duck Lagoon is managed by Kangaroo Island Council as a conservation reserve. The lagoon provides good habitat for flora and fauna, as illustrated by the presence of Australian White Ibis (*Threskiornis molucca*) and the Pied Cormorant (*Phalacrocorax varius*).

10.0 TENURE AND MANAGEMENT AUTHORITY

10.1 Background

The type of management authority and tenure surrounding a wetland often dictates the type and level of protection and management for the wetland. An understanding of this also allows consideration of different legislation and approaches concerning on-site management and planning for wetland sites.




Figure 10-1. Tenure and management

10.2 Analysis

The majority of wetlands (31 sites) are managed by landholders (eg Lashmar Lagoon, Rush Lagoon, Six-mile lagoon). Local government manages two wetland sites (Duck lagoon and Chapman River) and the Department for Environment and Heritage manages eight wetlands (eg Grassdale Lagoon, Bay Lake 2, Breakneck River). Note: includes 15 sites from information sourced from previous studies (Lloyd and Balla (1986) and Barritt (1982)).



Six mile lagoon 2 is private land. This lagoon is an excellent example of a freshwater wetland.

Plate 2. Six mile lagoon 2.

11.0 ENVIRONMENTAL ASSOCIATIONS AND IBRA REGIONS

11.1 Background

The Interim Biogeographic Regionalisation for Australia (IBRA) is a framework for conservation planning and sustainable resource management within a bioregional context. IBRA regions represent a landscape based approach to classifying the land surface from a range of continental data on environmental attributes. In 1999-2000, IBRA version 5 was developed. Eighty-five bioregions have been delineated, each reflecting a unifying set of major environmental influences that shape the occurrence of flora and fauna and their interaction with the physical environment. (See: http://www.ea.gov.au/parks/nrs/ibraimcr/ibra_95/index.html)

The Wetland Inventory for Kangaroo Island is located within the Flinders Lofty Block. The Flinders Lofty Block is described as temperate, with well-defined uplands of Cambrian and late Proterozoic marine sediments. The dominate vegetation type is eucalypt open forests and woodlands and heaths on mottled yellow and ironstone gravelly duplex soils in the wetter areas and red duplex soils in drier areas; now largely cleared for agriculture and urban development, (Thackway and Cresswell 1995).

Environmental associations are the next level of complexity down from the IBRA regions, these associations were first described and mapped by Laut *et al* (1977).

11.2 Analysis

The Wetland Inventory covers seven environmental associations, each having unique attributes. Refer to [Map 3](#) for Environmental Associations.




Figure 11-1. Number of wetlands located in Environmental Associations

Note: includes 15 sites from information sourced from previous studies (Lloyd and Balla (1986) and Barritt (1982).

The majority of wetlands surveyed are located in the Parndana environmental association, which covers the western and central areas of Kangaroo Island. Several wetlands located in this association have high conservation values, for example Larrikin Lagoon, Bay Lake and Six-mile Lagoon. The Mac Gillivray environmental association encompasses the saline lake district (eg Rush Lagoon, Wangara Swamp, White Lagoon and Salt Lagoon).

12.0 Key Biodiversity Areas

12.1 Background

Key biodiversity areas are outlined in the Biodiversity Plan for Kangaroo Island, developed by the Department for Environment and Heritage, South Australia. Key biodiversity areas have been developed through the use of existing biological data, vegetation mapping, GIS analysis, expert knowledge and community consultation (Willoughby, Opperman, Inns 2001). Three biodiversity areas are identified within the biodiversity plan; these are large remnant areas, threatened habitat areas and fragmented habitats. [Map 4](#) shows the location of these biodiversity areas.

Large remnant areas

These areas have been identified on the basis that they contain large blocks of vegetation, open space linkages, species diversity and populations of species with high conservation significance.

Threatened habitat areas

Threatened Habitat Areas have been identified on the basis that they are:

- selectively cleared and modified resulting in low remnancy of plant communities;
- contain vegetation that is poorly conserved within reserve systems;
- contain large numbers of species of high conservation significance.

Fragmented habitat areas

Fragmented habitats have been identified on the basis that they are:

- contain a mixture of large blocks of native vegetation and large amounts of cleared land;
- biodiversity in areas of large blocks likely to approach that of pre-European settlement levels;
- population sizes for most species are likely to be viable in the long-term.




Figure 12-1. Wetlands within Key Biodiversity Areas.

Note: includes 15 sites from information sourced from previous studies (Lloyd and Balla (1986) and Barritt (1982).

12.1 Analysis

The majority of wetlands surveyed are located within threatened habitats (22 sites) and includes wetlands surveyed in the saline district area and lagoons in Cygnet River. Wetlands surveyed in this region include Murray Lagoon, White lagoon, Salt Lake and Birchmore Lagoon. A large remnant biodiversity area is located on the western section of Kangaroo Island and encompasses Flinders Chase National Park. Thirteen wetlands were surveyed in this key biodiversity area. A biodiversity area of fragmented habitats is located in central Kangaroo Island and includes the upper drainage lines of several river systems; seven wetlands are identified in this area (eg Six-mile Lagoon).

13.0 WETLAND AREA

13.1 Background

The area of wetlands was calculated by using ArcView GIS based on waterbody mapping provided by Planning SA.

13.2 Analysis

The majority of wetlands have an area between 0 and 50 ha (18 sites). Four sites recorded areas between 51 and 100 ha and two sites recorded areas between 101 and 200 ha, namely Rush Lagoon 116 HA and Big White Salt Lagoon 105 ha. One site recorded an area between 201 and 300 ha (White Lagoon 291ha) and two sites recorded areas over 300 ha (Murray Lagoon 666 ha and Salt lagoon 400 ha). 27 wetlands were surveyed for this analysis.




Figure 13-1. Wetland area.

14.0 LANDFORM ELEMENT

14.1 Background

The landform element definitions used in the wetland inventory has been adapted from Heard and Channon (1997) *Guide to a native vegetation survey using the biological survey of South Australia methodology, Section 3. Geographic Analysis and Research Unit, Department of Housing and Urban Development*. For landform element descriptions, refer to [Appendix 1](#).

14.2 Analysis




Figure 14-1. Wetland landform elements.

Note: includes 15 sites from information sourced from previous studies (Lloyd and Balla (1986) and Barritt (1982)).

Wetlands are ultimately defined by the surrounding landforms. On Kangaroo Island, lagoons were the most common landform surveyed (14 sites). Examples of these lagoons include Duck Lagoon and Grassdale Lagoon. Salt lakes were also represented in the survey, and included wetlands located in the salt lake district located south of Kingscote.

15.0 GEOLOGY

15.1 Background

The geological record on Kangaroo Island spans nearly half of the Earth's history. The oldest exposed rocks (Cambrian 570 million years) on the island occur on the Dudley Peninsula the eastern end of the island. The basic framework of Kangaroo Island was formed during the Cambrian-Ordovician (505 million years ago). The younger sediments since then have formed the island's landscape. These sediments such as aeolianite dunes, spread across and along the south coast of Kanagroo Island altering the drainage flow from the higher central plataeu to the coast. This assisted in forming the lagoons and lakes in the southern section of Kangaroo Island, for example Murray Lagoon. Lunettes also formed during the late Pleistocene and are a unique feature on Kangaroo Island and good examples can be seen adjacent to White and Rush lagoons.

15.1 Analysis

Table 15-1. Geology underlying surveyed wetlands.

Geological description	Number of wetlands surveyed
Bridgewater formation	12
Alluvial/fluvial sediments	12
Kanmantoo trough	6
Lacustrine playa sediments	2
Gibbers, sandstones and claystones	6
Carboniferous permian rocks	2

Note: includes 13 sites from information sourced from previous studies (Lloyd and Balla (1986) and Barritt (1982).

The Bridgewater formation is located along the southern coast line and forms the underlying geology for 12 wetlands surveyed. These wetlands include; Grassdale Lagoon, Salt Lake and Big White Lagoon. Alluvial and fluvial sediments are found within another 12 wetlands surveyed, including, Murrays Lagoon, Rush Lagoon, White Lagoon and Salt Lagoon. Refer to [Map 5](#) for locations of geological formations underlying the surveyed wetlands.



The lunette composed of wind blown material and snail shells extend along the northern and eastern sides of the lagoon.

Plate 3. Lunette adjacent to White Lagoon.

16.0 HYDROLOGY

16.1 Background

The Wetland Inventory documented the annual average rainfall for the wetlands surveyed and made observations on the main sources of water entering the wetland. Many of the wetlands are closed hydrological systems being fed by small drainage lines and run-off from the surrounding catchment, a good example of this is Six mile Lagoon.

16.2 Analysis

Table 16-1. Rainfall analysis

Rainfall bracket	Number of wetlands
500	26
550	3
600	5
650	3
750	2
800	3

Note: includes 15 sites from information sourced from previous studies (Lloyd and Balla (1986) and Barritt (1982)).

Rainfall fall on the high plateau on the western section of the island contributes to surface water run-off into the south-eastern surrounding catchments containing wetlands. Wetlands not located in the south-eastern section of the island rely on localized surface water run-off to maintain the hydrological cycle.



Six-mile lagoon is an excellent example of a wetland dependent upon surface water entering the wetland from the surrounding catchment.

Plate 4. Six-mile Lagoon.

17.0 FLORA ANALYSIS

A large proportion of the wetlands surveyed on Kangaroo Island are considered to be freshwater (<1500 μ s). Aquatic and semi-aquatic flora was noticeably abundant in many water bodies, especially in freshwater (below 900 μ s). Members from the genera *Baumea*, *Blechnum*, *Carex*, *Gahnia* and *Juncus* are common and form the understorey component of the vegetation structure surrounding these wetlands. Saline waterbodies recorded aquatic flora such as *Samolus repens*, *Ruppia*, *Chara*, *Wilsonia* and *Myriophyllum* species. These genera were common within White lagoon, Rush Lagoon and Murray Lagoon.

Table 17-1. Vegetation associations at survey sites

Dominant vegetation community adjacent to wetland	Number of wetland sites
<i>Melaleuca halmaturorum</i> over introduced grasses	7
<i>Melaleuca halmaturorum</i> tall shrubland over <i>Gahnia filum</i> open sedgeland	3
<i>Melaleuca halmaturorum</i> tall shrubland over <i>Sclerostrogia arbuscula</i> Tall open shrubland	8
<i>Eucalyptus diversifolia</i> , <i>E. cosmophylla</i> , +/- <i>E. baxteri</i> , +/- <i>E. fasciculosa</i> +/- <i>Melaleuca gibbosa</i> over <i>Gahnia</i> sp. Open mallee.	1
<i>Eucalyptus camaldulensis</i> var. <i>camaldulensis</i> , <i>E. leucoxydon</i> ssp. <i>leucoxydon</i> over introduced grasses	2
<i>Eucalyptus cladocalyx</i> , <i>E. diversifolia</i> , +/- over <i>Callistemon rugulosus</i> +/- <i>Juncus</i> sp. woodland	1
<i>Eucalyptus remota</i> , <i>E. cosmophylla</i> , +/- <i>E. baxteri</i> over <i>Acacia retinoides</i> and <i>Gahnia</i> sp	1
<i>Eucalyptus cosmophylla</i> , <i>E. baxteri</i> over <i>Leptospermum continentale</i> , <i>Acacia retinoides</i> var. <i>retinoides</i> , Low woodland	2
<i>Eucalyptus baxteri</i> , <i>E. obliqua</i> , <i>E. cosmophylla</i> low woodland	1
<i>Melaleuca gibbosa</i> , <i>M. brevifolia</i> +/- <i>Callistemon rugulosus</i> Var. <i>rugulosus</i> +/- <i>Hakea rugosa</i> shrubland	2
<i>Eucalyptus cneorifolia</i> , <i>E. lansdowneana</i> ssp. <i>albopurpurea</i> , <i>E. diversifolia</i> , +/- <i>E. cosmophylla</i> Open mallee	1
<i>Melaleuca brevifolia</i> , <i>M. gibbosa</i> , +/- <i>M. halmaturorum</i> Shrubland	3
<i>Eucalyptus cladocalyx</i> , +/- <i>E. obliqua</i> , +/- <i>E. baxteri</i> , <i>E. cosmophylla</i> Woodland	1
<i>Eucalyptus cosmophylla</i> , <i>Eucalyptus fasciculosa</i> , very open mallee +/- <i>Melaleuca halmaturorum</i> over <i>cyperus</i> sp	1
<i>Leptospermum continentale</i> , <i>A. retinodes</i> var. <i>retinodes</i> , <i>E. cosmophylla</i> , <i>E. baxteri</i> Shrubland.	1

Note: includes eight sites from information sourced from previous studies (Lloyd and Balla (1986) and Barritt (1982).

Willoughby *et al* (2001) state that vegetation associations containing *Melaleuca halmaturorum*, *Melaleuca brevifolia* and *Melaleuca gibbosa* are considered to be rare on Kangaroo Island. The association of *Callstemon rugulosus* var. *rugulosus*, +/- *Melaleuca halmaturorum*, +/- *Hakea rugosa* Tall closed shrubland is also considered to be rare on Kangaroo Island.

The ecological role of *Melaleuca halmaturorum* as a fringing saline wetland species is very important. *Melaleuca halmaturorum* often forms an effective buffer for water bodies from increased sediment loads and nutrient concentrations. Greenway (1997) discusses some further direct benefits from *Melaleuca halmaturorum* these include:

Hydrological benefits

- Improved water quality by filtering suspended particles and by removing, recycling, or immobilising contaminants and nutrients, thereby preventing deterioration of downstream aquatic ecosystems (eg Cygnet River Lagoon).
- Provide a protective buffer zone between shorelines, estuaries and river systems protecting these waterways from siltation, nutrient run-off and erosion
- Provide flood mitigation by storing and detaining precipitation and run-off thus reducing flow rates and peak floods (eg Rush and White lagoons).
- Provide groundwater recharge and a water source for people and wildlife.

Ecological benefits

- *Melaleuca* trees are highly productive at recycling nutrients and function as long-term biomass sinks (eg Destree Swamp).
- During major flood events, particulate matter is washed into the rivers and estuaries to provide a food source for heterotrophic micro-organisms and detritivores (eg Lashmar Lagoon).
- Provide both temporary and permanent habitats for a variety of flora and fauna, including roosting and breeding areas for wildlife. Some *Melaleuca* swamps support large ibis and egret colonies.
- Provide vital refuges for wildlife during periods of drought.
- *Melaleuca* trees flower prolifically and provide a source of nectar for resident and migratory birds, bats, possums, bees and other insects. Their nectar is a particularly valuable food source for migratory honey-eaters and parrots during the autumn/winter months.

Threats

There are various key threatening processes which affect the majority of remnant *Melaleuca halmaturorum* woodlands on Kangaroo Island. These include increased nutrients moving into the water due to catchment clearance and farming practices, resulting in dieback and a change in water chemistry.



Plate 5. Halls Road Swamp

Halls Road Swamp has suffered from increased salinity levels and reduced frequency of freshwater flushing which has caused extensive dieback of the *Melaleuca halmaturorum*



Plate 6. Destree Swamp

Destree Swamp located on the Hundred Line Road is an excellent example of a functioning Melaleuca swamp.



Plate 7. Cygnet River Lagoon

Cygnet River Lagoon plays an important role in filtering nutrients before water enters Cygnet River.

18.0 DEGRADATION AND DISTURBANCE

18.1 Background

Disturbances or threats are defined as any direct or indirect human activities at the site or in the catchment area that may have a detrimental effect on the ecological character of the wetland. The effect may be a low level disturbance (low level grazing) or a major threat such as water diversion schemes.

18.2 Analysis

Table 18-1. Land degradation

Degradation type	Number of Occurrences
Access Tracks	13
Clearance	22
Grazing damage	14
Fence lines	5
Mining impact	1
Altered flows	3

Many wetlands contained disturbances; the two most frequent disturbances recorded are clearance (22 occurrences) and grazing damage (14 occurrences). Access tracks dissecting or traversing wetlands were also common (13 sites). A concern with many of these tracks is that they are usually formed with compressed soil and during high rainfall events, sediment often runs into adjacent wetlands. Other disturbances recorded include fences (5 sites) and altered water flows (3 sites) through stormwater inputs, banks, drainage and water extraction.

Other disturbances not recorded in this survey but documented by other observers, include the impacts from feral pigs (*Sus scrofa*) and populations of introduced freshwater crayfish, Yabbies (*Cherax destructor*) and Marron (*Cherax tenuimanus*) (Willoughby 2001).



An example of a common land degradation disturbance. Clearance is of a major concern and is affecting many wetland ecosystems on Kangaroo Island. Increased salinity levels through vegetation clearance can destroy freshwater ecosystems.

Plate 8. Halls Road Swamp

19 AQUATIC VEGETATION CLASSES

19.1 Background

Parameters for seven classes of aquatic vegetation were included in the survey. These records can indicate the types of producers (production of oxygen and plant food) within the wetland system. The diversity of classes recorded may indicate the level of aquatic biodiversity present, the vegetation classes consist of:

Algal and aquatic moss commonly comprise Charophyta (stoneworts) and Chlorophyta (green algae) which forms macroscopic mats either attached to plants or in open water. Algae forms the photosynthetic basis for the open water food sources in many inland waters, (Boulton and Brock, 1999).


For images of green algae see: <http://www.nmnh.si.edu/botany/projects/algae/Imag-Chl.htm>.

Floating vascular/leaved plants have part or all of the leaves at the waters surface. Examples include *Azolla* species floating ferns that host bacteria that fix nitrogen, (Romanowski 1998), *Lemna*, *Spirodela* and *Wolffia* (duckweeds) and the genera *Utricularia* (bladderworts). Members from the family Potamogetonacea (pondweeds) are also common floating plants and can be found in a variety of habitats. All these plants are able to provide habitat for invertebrates, provide shelter for fishes and produce oxygen.

Rooted vascular plants are those rooted in the sediments with either a major proportion of material above water (reeds, rushes and sedges) or totally under water (*Vallisneria* spp.). Many of these plants play a key role in nutrient cycling and provide habitat for birds, insects and aquatic invertebrates. Typical genera include *Baumea*, *Bolboschoenus*, *Carex*, *Cyperus*, *Gahnia*, *Schoenus*, *Juncus*, *Triglochin* and *Myriophyllum*. *Myriophyllum* is a distinctive wetland genus that provides food, shelter and spawning or nesting sites for a variety of animals, from invertebrates to fish, frogs and birds (Romanowski, 1998).

19.2 Analysis

Figure 19-1. Aquatic vegetation types



Note: includes eight sites from information sourced from previous studies (Lloyd and Balla (1986) and Barritt (1982).

Rooted vascular plants are the most common form of vegetation class within the surveyed wetlands. Genera such as *Juncus*, *Carex*, *Cyperus* and *Gahnia* were frequently recorded in freshwater wetlands (29 sites, eg Grassdale Lagoon). Saline wetlands commonly recorded species taxa such as *Samolus repens* (creeping brookweed), *Ruppia* ssp and *Myriophyllum* spp (eg Rush Lagoon).



The distinct bands of floating vegetation can be seen in the center of the photo.

Plate 9. Rush Lagoon



Grassdale lagoon has a diversity of all aquatic classes. Pictured are dense stands of Juncus.

Plate 10. Grassdale Lagoon



Discovery Lagoon recorded a high abundance and diversity of submerged aquatic flora, including Chara and Myriophyllum spp.

Plate 11. Discovery lagoon

20 AQUATIC FAUNA ANALYSIS

20.1 Invertebrates

Macro and micro invertebrates are an essential component of the wetland food web. They are responsible for a significant proportion of the secondary production occurring in wetlands, and form two interconnected wetland food chains, a grazing food chain and a detrital food chain. Invertebrates comprise much of the diet of waterfowl populations the diversity and abundance of waterfowl can be a direct consequence of the invertebrate food supply.

20.1.1 Ecological benefits

Yen and Butcher (1997) provide some examples of direct ecological benefits that invertebrates contribute:

Tangible direct benefits:

1. Plant pollination
2. Effects on soil; soil formation and fertility.
3. Decomposition; fragmentation and recycling of dead plant and animal material.
4. Position in the food web; invertebrates are the principle food for many vertebrates.
5. Predation and parasitism. Invertebrates are involved in the natural regulation of populations of other species through predation and parasitism, and thus form the basis of biological control.

Indirect ecological benefits:

1. Ecosystem stability: the loss of species from highly interrelated systems is likely to cause a cascade of further extinction losses.
2. Evolutionary time: the diversity within ecosystems over time maintains diversity.

20.1.2 Trophic dynamics

Standing water communities are dynamic systems which reflect change in many variables. The trophic state of a wetland depends on nutrient inputs from the catchment and within the wetland (Boulton and Brock 1999). Invertebrates were collected during the Wetland Inventory and classified according to trophic levels. If samples from all trophic groups are collected, this could suggest that the aquatic ecosystem is in a reasonable state of equilibrium. The top of the food chain is occupied by vertebrate predators, including fish, water rats and water birds. These terrestrial predators can be considered to be on the top of the aquatic food chain, and provide a pathway for the export of nutrients and other material from the wetland ecosystem (Boulton and Brock, 1999). These trophic levels are described below.

Primary producers

Primary producers form two groups, those that are suspended or floating and those attached to substrate or other plants. Attached macrophytes includes fringing reeds and submerged plants and periphyton (the biota attached to submerged surfaces). Suspended or floating forms generally consist of the phytoplankton and algae groups. Phytoplankton form the basic photosynthetic basis for the open water food web in most standing waters.

Consumers

There are two main types of consumers based on diet: grazers that consume plants and predators that consume other animals.

Grazers

Grazers consist of aquatic snails (Gastropoda) and some mayfly nymphs (Ephemeroptera), caddisfly larvae (Trichoptera) and beetles (Coleoptera). These groups are usually found near the edges of the water body.

Within the open water, some of the important grazers are zooplankton, including water fleas (Cladocera) and copepods (Calanoida and Cyclopoida).

Vertebrate Grazers

Vertebrate grazers generally consist of groups such as tadpoles, fish and waterbirds. Vertebrate grazers can influence the food web considerably when attracted to water bodies in times of flood or in types of drought.

Predators

Predators include dragonfly larvae (Odonata) which tend to ambush prey and invertebrates that hunt in open water such as diving beetles (Dytiscidae, Coleoptera) (Boulton and Brock 1999). Areas such as the littoral zone tend to have high biodiversity of grazers which in turn attracts many invertebrate predators.

20.2 Temporary Wetlands

Many of the wetlands on Kangaroo Island are temporary and display slightly different invertebrate fauna composition from other wetland systems such as saline lakes or permanent waters. Williams (2000) makes four general conclusions from his study of temporary wetlands, these are:

1. Faunal diversity is high and often higher than in many permanent wetlands.
2. A wide range of fauna groups occurs with the particular assemblage depending largely upon time from filling. Many species are restricted to temporary wetlands, for example all notostracan, conchostracan and anostracan species are restricted to temporary wetlands.
3. Local differences in hydrology, filling frequency, basin shape and other factors often result in differences between wetlands in the same area and same time.
4. Considerable continental and regional endemism prevails. Most macrofaunal species are endemic to Australia.

The filling or flooding of temporary water bodies releases a pulse of nutrients that, together with light and water, provide the resources for germination and growth of both micro and macro photosynthesizers. Habitat for consumers and decomposers soon follows. The invertebrate sediment egg bank with desiccation-resistant stages seems to be the initial source of colonists. The groups that tend to be first in temporary waters include rotifers, ostracods, copepods and cladocerans. When the water body starts to dry a 'predator soup' results, and terrestrial predators (eg birds) come to the water to feed during the drying process. This process forms a critical trophic link between aquatic and terrestrial systems (Boulton and Brock 1999).

20.3 Saline Systems

Large areas of Kangaroo Island contain saline wetlands, those surveyed include:

- Lashmar Lagoon
- Greenfields Wetland
- Big White Salt Lagoon
- Discovery Lagoon
- Salt Lake
- Wangara Lagoon
- Destree Swamp
- Murray Lagoon
- Chapman River
- Salt Lake 2
- Milky's Lake
- Rush Lagoon
- White Lagoon 1
- Salt Lagoon
- Halls Road Swamp
- Emanual Lagoon
- White Lagoon
- Birchmore Lagoon
- Nepean Bay Lagoon

Freshwater organisms in Australia generally tolerate salinities up to about 3 g/L (3000 EC), beyond this there is a change in community composition, with decreased richness and increased abundance (Williams, 1998); Skinner *et al* 2001). The biological process in salt lakes can resemble those of fresh water bodies despite the differences in physical, chemical and biological attributes (Boulton and Brock 1999). The beds of many salt lakes are covered by benthic microbial mats dominated by photosynthetic producers, and lake crusts contain propagules of decomposers, producers and consumers. Boulton and Brock (1999), comment that little is known about the microbial loop in salt lakes.

A high diversity of invertebrates can occur within salt lakes, examples include rotifers, anostracan, cladocerans, calanoid copepods and ostracods. Fish are usually absent from saline lake systems and the top consumers are mostly water birds.

In general, invertebrate species richness in salt lakes declines with increasing salinity, but at intermediate salinities where many species tolerances are broad, other factors such as biological interactions and pH will affect community composition (Skinner *et al* 2001, Williams, 2000). Studies by Skinner *et al* (2001) indicates that salinization shifts invertebrate community structure and algae tends to also become dominant at the higher salinity levels. This could lead to insufficient food for animals higher in the trophic level, including fish and waterfowl.

20.4 Analysis

Analysis of invertebrates is discussed in four areas:

1. frequency of invertebrate occurrence
2. trophic levels
3. number of invertebrate records for each surveyed wetland
4. invertebrate lists for each wetland tabled in Appendix 2.

20.4.1 Frequency of invertebrate occurrence

Table 20-1 lists the invertebrate species identified from collections made during the Wetland Inventory in December 2000, with the frequency of occurrence given against each identified invertebrate. A total of 80 species are recorded, with 37 species recorded once and 43 species recorded more than once. Of these 43 species, three species recorded 10 or more occurrences, (*Austrochiltonia australis* 10, *Diacypris spinosa*, 13 and *Mesocyclops sp.* 12 occurrences).



Plate 12. *Diacypris* cf. *spinosa*.

The genera *Diacypris* (recorded 13 times) belongs to the subclass Ostracoda, of which there are 11 families and 37 genera.

Ostracods are generally found in most habitats Australia-wide.



Plate 13. *Dicrotendipes* sp.

Five occurrences of *Dicrotendipes* sp. were recorded.



Plate 14. *Tricoptera oecetis*.

Two occurrences of *Tricoptera oecetis* (caddis fly) were recorded. Note the protective case surrounding the larvae.

Table 20-1. Frequency of invertebrate species occurrence

Identified invertebrates	Frequency of occurrence
<i>Ablabesmyia</i> sp	1
<i>Aeshnia brevistyia</i>	1
<i>Agraptocorixa</i> sp	3
<i>Alboa worooa</i>	6
<i>Alona</i> sp	2
<i>Anisops</i> sp	8
<i>Apocyclops annulosus</i>	2
<i>Australocyclops australis</i>	5
<i>Australocyclops similis</i>	2
<i>Australocyclops</i> sp	1
<i>Austrochiltonia australis</i>	10
<i>Austrolestes annulosa</i>	2
<i>Austrolestes cingulatus</i>	2
<i>Austrolestes leda</i>	1
<i>Austrolestes</i> sp	2
<i>Berosus</i> sp.	2
<i>Boeckella</i> sp	2
<i>Boeckella symmetrica</i>	7
<i>Boeckella triarticulata</i>	2
<i>Calamoecia ampulla</i>	5
<i>Calamoecia australis</i>	1
<i>Calamoecia ciittelata</i>	3
<i>Calamoecia salina</i>	4
<i>Candonocypris</i> sp	2
<i>Ceriodaphnia</i> sp	1
<i>Chironominus</i> sp	6
<i>Cladotanytarsus</i> sp	1
<i>Chostonectes</i> sp	1
<i>Cloeon</i> sp	1
<i>Cnephia</i> sp	1
<i>Culex</i> sp	2
<i>Cypricercus</i> sp	1
<i>Daphnia cf carinata</i>	1
<i>Daphnia</i> sp	1
<i>Daphniopsis</i> sp	2
<i>Diacyclops bisetosus</i>	1
<i>Diacypris spinosa</i>	13
<i>Dicrotendipes</i> sp	5
<i>Dunhevedia crassa</i>	2
<i>Enithares</i> sp	4
<i>Enithares woodwardi</i>	2
<i>Eucyclops</i> sp	1
<i>Eucyclops cf baylyi</i>	1
<i>Fluvidona</i> sp	1
<i>Gomphodella</i> sp	1
<i>Haloniscus searli</i>	1
<i>Hemianax papuensis</i>	1
<i>Hesperocordula</i> sp	1
<i>Ilyocryptus</i> sp	2
<i>Kennethia</i> sp	2
<i>Keratella</i> sp.	1
<i>Limnocythere porhyretica</i>	1
<i>Macrothrix</i> sp	9
<i>Macroscyclops</i> sp	1
<i>Mesocyclops</i> sp.	12
<i>Metacyclops mortoni</i>	6
<i>Miconecta</i> sp	8
<i>Mixocyclops</i> sp	2
<i>Mytilocypris tasmanica</i>	1
<i>Mytilocypris ambigua</i>	4
<i>Necterosoma</i> sp	1
<i>Neocyclops petkovis</i>	1
<i>Newnhamia fenestrata</i>	2
<i>Notalina</i> sp	1
<i>Odontomyia</i> sp	1
<i>Oxythira cf columba</i>	1
<i>Paracyclops</i> sp	1
<i>Parachironomus</i> sp	1
<i>Parartemia zeitiana</i>	7
<i>Paraterina</i>	1
<i>Paratanytarsus</i> sp	1
<i>Plea</i> sp	1
<i>Procladius</i> sp	5
<i>Sigara</i> sp	2
<i>Simocephalus cf expinosus</i>	1
<i>Simocephalus elizabethae</i>	2
<i>Simocephalus</i> sp.	3
<i>Tanytarsus</i> sp	6
<i>Tricoptera oecetis</i>	2
<i>Triplectides australis</i>	1

Invertebrate identification by Robert Walsh, 2000.

20.4.2 Trophic levels

Invertebrate composition analysis is affected by the study of trophic levels present in each wetland. Conclusions made about the ecosystem health of a wetland based on the invertebrates could not be made due to the limited sample size, and can only provide an indication at the time of the sampling.

Invertebrate scoring method

Invertebrate scores were assigned for each wetland, each sample was identified to family level and the trophic level of each family recorded. From the samples identified the diversity and abundance was recorded and the following scoring system has developed.

Score 1 = (Low) sampled one family from one trophic level, usually from a lower level trophic level, (eg detritivores and herbivores).

Score 3 = (Moderate) sampled more than one family with representatives from one or more trophic levels, (eg herbivores and carnivores).

Score 5 = (High) sampled more than two families with representatives from three or more trophic levels, (eg detritivores, herbivores, omnivores and carnivores).




Figure 20-1. Invertebrate scores.

Table 20-2. Trophic scores

Wetland number	Wetland name	Trophic score
S5000	Lashmar Lagoon	5
S5006	Milkys Lake	5
S5009	Moores Lake	No data
S5013	Greenfields Wetland	3
S5018	Rush Lagoon	5
S5020	Big White Salt Lagoon	1
S5021	White Lagoon1	5
S5030	Discovery Lagoon	1
S5031	Salt Lagoon	3
S5035	Salt Lake	3
S5041	Halls Road Swamp	5
S5042	Wangara Lagoon	3
S5043	Emanual Lagoon	5
S5044	Destree Swamp	3
S5050	White Lagoon	5
S5053	Murray Lagoon	5
S5055	Birchmore Lagoon	No data
S5056	Six Mile Lagoon	5
S5061	Chapman River	No data
S5062	Nepean Bay Lagoon	3
S5063	Salt Lake 2	3
S5064	Duck Lagoon	5
S5065	Cygnets River Lagoon	5
S5066	Six Mile Lagoon 2	5
S5069	Bay Lake 2	No data
S5070	Breakneck River	5
S5071	Grassdale Lagoon	5

*Note four sites had no records



Chironomidae, 40X
(non-biting midges,
bloodworms).

Chironomidae
have been
recorded in many
wetlands on
Kangaroo Island,
such as Grassdale
Lagoon and Rush
Lagoon.

They are found
Australia-wide in
all habitats, and fill
the trophic level of
omnivores.

Plate 15. Chironomidae



The genus *Anisops*
is an active
predator.

This genera was
recorded in several
wetlands, (eg Six-
mile Lagoon and
Cygnet River
Lagoon).

Plate 16. Anisops.

20.4.3 Invertebrate records for each surveyed site

Table 20-3 lists the number of invertebrate records identified at each wetland site surveyed. Wetlands with notable records include Rush Lagoon (36 records), Duck Lagoon (33 records), Grassdale Lagoon (27 records) and Six-mile Lagoon 2 (24 records).

Table 20-3. Number of invertebrate records for wetlands

Wetland number	Wetland name	Number of records
S5000	Lashmar Lagoon	5
S5006	Milkys Lake	6
S5009	Moores Lake	No data
S5013	Greenfields Wetland	13
S5018	Rush Lagoon	36
S5020	Big White Salt Lagoon	1
S5021	White Lagoon1	6
S5030	Discovery Lagoon	2
S5031	Salt lagoon	9
S5035	Salt Lake	2
S5041	Halls Road Swamp	9
S5042	Wangara Lagoon	4
S5043	Emanual lagoon	7
S5044	Destree Swamp	4
S5050	White Lagoon	12
S5053	Murray Lagoon	6
S5055	Birchmore Lagoon	8
S5056	Six Mile Lagoon	15
S5061	Chapman River	No data
S5062	Nepean Bay Lagoon	3
S5063	Salt Lake 2	5
S5064	Duck Lagoon	33
S5065	Cygnnet River Lagoon	14
S5066	Six Mile Lagoon 2	24
S5069	Bay Lake 2	No data
S5070	Breakneck River	15
S5071	Grassdale Lagoon	27

20.5 Amphibians

No amphibians were observed during the wetland inventory survey, however past studies have commented on amphibians within wetlands on Kangaroo Island. Willoughby *et al* (2001) comments that six frog species have been recorded on Kangaroo Island. One of these is the Brown toadlet (*Pseudophryne bibroni*), the status of which is largely unknown.

20.6 Fish

Fish were observed in high abundances within the Chapman River. These species were probably *Galaxias maculatus* and *Atherinosoma microstoma*. No other fish species were observed during the wetland inventory.

20.7 Mammals

No mammals of conservation significance were observed during the survey. However, Neagle (2002) has identified the Swamp Rat (*Rattus lutreolus*) as a species in severe decline. Lloyd and Balla (1986) recorded the Swamp Rat at Grassdale Lagoon during their 1986 study.

20.8 Avi-fauna

No species of conservation significance were identified during the survey, but species such as the Musk Duck (*Biziura lobata*) which is listed as rare at a State level, has been recorded at Murrays Lagoon (Lloyd and Balla, 1986). The wetland inventory did record abundant bird activity on many lagoons and lakes but some of the more common species observed include:

- Pacific Black Duck (*Anus superciliosa*)
- Masked Lapwing (*Vanellus miles*)
- Black Swan (*Cygnus atratus*)
- White Faced Heron (*Egretta novaehollandie*)
- Eurasian Coot (*Fulica atra*)
- Banded Lapwing (*Vanellus tricolor*)
- Australian Shellsuck (*Tadorna tadornoides*)
- Black Winged Stilt (*Himantopus himantopus*)

21 WATER CHEMISTRY

Chemical processes in permanent and temporary waters are extremely complex. The chemistry of the water directly influences the biological process (such as photosynthesis). The physical features of the wetland also has a strong influence on both the chemical and biological processes. These three factors (chemical, physical and biological) are constantly in a state of flux and change. Changes in these parameters are pronounced in temporary wetlands where a wetting and drying cycle occurs. The majority of wetlands within Kangaroo Island have seasonal water regimes, filling during winter and remaining dry throughout summer.




Figure 21-1. Temporary wetland cycles

Brock and Boulton (1999) state that changes in water quality during drying and filling depend on factors, including:

1. sediment properties (composition, nutrients and organic content)
2. type of drawdown (gravity or evaporative)
3. severity of drying (rate of drying, temperature, weathering)
4. conditions of refilling (origin of water, degree of sediment disruption).

Figure 21-1 shows the changes in chemical variables over time during the phases of filling and drying in temporary wetlands. Seasonal changes in invertebrate composition is also noted. (Adapted from Boulton and Brock, 1999).

21.1 pH

21.1.1 Background

The pH value of water indicates how acidic or alkaline it is on a scale 1-14. Acids have a low pH of about 2 for a strong acid like sulphuric acid and about 4 for a weak one like lactic acid. Alkalis have a high pH of about 12 for sodium hydroxide. Pure distilled water has a pH of 7 which is neutral. From pH 7 to 0, a liquid becomes increasingly acidic and from pH 7 to 14, a liquid becomes increasingly alkaline.

Generally in South Australia, the pH of natural water ranges between 6.0 and 8.5 with most water bodies in the range 7.0-8.0. The higher pH of natural water bodies are caused by the high bicarbonate levels in the water and can raise the pH during the day and lower pH at night. Agricultural chemicals entering the water can also affect the pH.

pH is an important environmental indicator. At extremely high or low pH values, the water becomes unsuitable for most organisms.

14	HIGH (Alkaline)
10	MEDIUM
9	MILD
8	PRISTINE
6	MILD
5	MEDIUM
4	HIGH (Acidic)
1	

Figure 21-2. pH values

21.1.2 Analysis

The average pH of wetlands surveyed ranged between pH 6 and 9 (19 sites), this falls into a neutral pH value. Six sites recorded a pH reading of alkaline (pH between 9-14), Rush Lagoon and Halls Road Swamp received the highest reading of 9.7. The other alkaline wetlands include Discovery Lagoon, Destree Swamp, White Lagoon and Birchmore Lagoon.

21.2 Conductivity

21.2.1 Background

Salinity is a serious threat to aquatic ecosystems throughout Australia. The National Land and Water Resources Audit 2001 estimates that up to 41,300 kilometers of streams could be salt affected by 2050 and that 24 of 79 river basins studies exceeded tolerable salinity parameters.

Salinisation causes serious biological effects, including;

- changes to the natural character of water-bodies
- loss of biodiversity
- less salt-tolerant species are replaced by more tolerant species.

These effects can cause permanent degradation and ecosystem collapse (Williams 2001). The loss of biodiversity is probably greater than generally realised since very little research has occurred in this area. Salinisation also leads to significant decreases in water quality for irrigation and water supply (leading to high economic costs); the loss of amenity and aesthetic values is also of concern.

Limited data are available for assessing the risk of adverse effects from salinity in different ecosystem types, particularly wetlands. The Australian and New Zealand Guidelines for Fresh and Marine Water Quality (2001) tables wetlands in South Australia with 'no data' related for assessing adverse effects of salinity. Guidelines developed for Western Australia state that lakes, reservoirs and wetlands should have a salinity range between 300-1500 μ s. Values at the lower end of the range are during rainfall events, those at the higher range are found in salt lakes and marshes.

Figure 21-1 is based on the salt measurement conversions guide from the Land Management Society Inc. This guide indicates salinity levels and tolerance levels for different parameters. These guidelines form the basis of analysis for the Wetland Inventory.

21.1.2 Analysis

Table 21-1 outlines the salinity ranges into which the wetlands fell. Salinity is one of the major threats to the freshwater ecology of wetlands on Kangaroo Island. Those wetlands that recorded low conductivity readings such as Six-mile Lagoon, should be monitored to assess the likely impacts that salinity will have on such systems. The naturally saline salt lakes are also suffering from increasing conductivity, with levels reaching saturation point. At these high levels aquatic flora and fauna cease to exist.

Table 21-1. Salinity guidelines.

Salinity in micro-siemens per cm $\mu\text{s/cm}$	Water definition	Guides	Surveyed Wetlands
0 - 900	Fresh	Fresh	Breakneck River, Six-mile Lagoon
900 - 2700	Marginal	Maximum for hot water systems, dam water starts to go clear, maximum for people.	Cygnets River Lagoon, Six-mile Lagoon 2, Grassdale Lagoon
3000 - 9100	Brackish	Maximum for milking cows and poultry, crop losses start.	Duck Lagoon
9500 +	Salt	9.5 = Yabbie growth starts to slow, maximum for horses. 14.5 = Yabbie growth ceases	Lashmar Lagoon, Milky's Lake, Greenfields Wetland, Rush Lagoon, Big White Salt Lagoon, White Lagoon 1, Discovery Lagoon, Salt Lagoon, Salt Lake, Halls Road Swamp, Wangara Lagoon, Emanuel Lagoon, Destree Swamp, White Lagoon, Murray Lagoon, Birchmore Lagoon, Chapman River, Nepean bay Lagoon, Salt Lake 2.



Six-mile Lagoon recorded the lowest conductivity reading (3.15 μs)

Plate 17. Six-mile Lagoon.



Duck Lagoon recorded a brackish conductivity reading of 7.70 mS/cm.

Plate 18. Duck Lagoon



Big white salt lagoon recorded a conductivity reading of 134.5 mS/cm.

Plate 19. Big White Salt Lagoon.



'Salt lake', a naturally saline lake recorded a reading of 188.8 mS/cm. Note the calcrete shoreline, this area was once used for gypsum mining.

Plate 20. Salt Lake.

21.3 Turbidity

Turbidity is a measure of water clarity and can be affected by the amount of suspended particles of clay, silt, plankton, industrial wastes and sewage in water. Turbidity is caused by small particles suspended in water which are too small to settle out, but big enough to scatter light. At higher levels of turbidity, water loses its ability to support a diversity of aquatic organisms, and water becomes warmer as suspended particles absorb heat from the sun. Turbidity leads to less light penetrating the water, thereby decreasing photosynthesis, which in turn reduces dissolved oxygen concentrations. Suspended particles may also clog aquatic invertebrate appendages and fish gills, reduce growth rates, and prevent egg and larval development. The plant community structure and biomass in an ecosystem is determined by the light regime, in conjunction with nutrients and temperature. Changes in water clarity may produce changes in the dominant phytoplankton groups. A common result from increased clarity tends to reduce the primary productivity of systems. Changes to the primary producers will also effect the other trophic components of the ecosystem (Liston and Maher, 1997).

Turbidity can have a significant effect on the microbiological quality of drinking-water, its presence can interfere with the detection of bacteria in drinking-water. Turbid water has been shown to stimulate bacterial growth since nutrients are adsorbed on to particulate surfaces, thereby enabling the attached bacteria to grow more rapidly than those in free suspension. The major problem associated with turbidity is its effect on disinfection. High levels have been shown to protect microorganisms from the action of disinfectants and to increase the chlorine and oxygen demand.

(http://www.who.int/water_sanitation_health/GDWQ/Chemicals/turbidity.htm).

Turbidity in water is measured in units called Nephelometric Turbidity Units (NTU's). It is calculated by measuring the dispersion of a light beam passed through a sample of water. Fine particles, silt, and suspended matter will cause a light beam passing through the water to be scattered. It has been found that the amount of scattering is proportionate to the amount of turbidity present. Therefore, this process gives a good indication of the relative turbidity of a water sample.

The measurement of NTU does not give the sizes of the particles, nor does it indicate the amount of particles present. It is a qualitative, rather than quantitative way of measuring turbidity.

21.3.1 Background

To understand the results fully, information about the natural levels of turbidity within all the survey wetlands at various times of the year is really required. This is important because normal levels of turbidity can vary greatly in wetlands from clear flowing zones to murky areas. Natural variations are also related to flow events such as floods, winter rains, road run-off which can increase levels dramatically.

The *Australian Water Quality Guidelines for Fresh and Marine Waters* do not set a turbidity guideline for the protection of aquatic ecosystems, but recommend that increases in suspended solids should be less than 10% of the seasonal mean NTU. This reinforces the need to ascertain the natural turbidity levels within waterbodies in Kangaroo Island. Bek and Robinson (1991) also suggest that turbidity below 50 NTU is suitable for protecting aquatic animals and plants. The 2000 update of *Australian Guidelines for Water Quality Monitoring* does provide some turbidity ranges for South-Central Australia. These ranges however are so broad they cannot be used as a measure against the Wetland Inventory turbidity results for Kangaroo Island.

The New South Wales Environmental Protection Agency, (EPA) has developed the following general guidelines for turbidity, these guidelines provide a useful method to compare the turbidity values collected during the Wetland Inventory.

- good (< 5 NTU)
- fair (5-50 NTU)
- poor (> 50 NTU)

Refer to: <http://www.epa.nsw.gov.au/soe/97/ch3/11.htm> for more information.

The National Advisory Committee to Water on the Web has developed regional trends of fresh water fish activity against turbidity values and time, (figure 21-5).

(See: <http://wow.nrri.umn.edu/wow/under/parameters/turbidity.html> for more information)

These trends correspond well to those guidelines developed by the NSW EPA, as mentioned above.




Figure 21-3. Fish activity against turbidity values and time

21.3.2 Analysis

Table 21-2 illustrates the number of surveyed wetlands against the turbidity guidelines developed by the NSW EPA. The majority of wetlands recorded a NTU value less than 5 (17 sites). Seven sites recorded NTU readings between 5-50, and Big white salt lagoon recorded the highest reading of 326 NTU. This wetland has an abundance of *Parartemia zeitziana* (fairy shrimps) which has agitated the fine sediments resulting in high turbidity.

Guideline	NTUs	Number of Wetlands
Good	<5	17
Fair	5-50	7
Poor	>50	1

Table 21-2. Turbidity guidelines.

21.4 Water Temperature

21.4.1 Background

Many of the physical, biological and chemical characteristics of a wetland are directly affected by temperature. Temperature affects the solubility of oxygen in water with warmer water holding less oxygen than cooler water. These factors affect the rate of photosynthesis by algae and larger water plants, with warm water being susceptible to eutrophication and therefore algal blooms. The sensitivity of organisms to toxic waste, parasites and diseases is also related to water temperature. Organisms can become stressed as temperatures rise and become less resilient to other stresses. This results in most aquatic organisms having a narrow temperature range in which they can function effectively.

Most aquatic organisms are cold-blooded which means they are unable to internally regulate their core body temperature. Therefore, temperature exerts a major influence on the biological activity and growth of aquatic organisms. To a point, the higher the water temperature, the greater the biological activity. Fish, insects, zooplankton and other aquatic species all have preferred temperature ranges. As temperatures get too far above or below this preferred range, the number of individuals of the species decreases until finally there are few, or none.

The most obvious reason for temperature change in lakes is the change in seasonal air temperature. Daily variation also may occur, especially in the surface layers, which are warm during the day and cool at night. In deeper lakes (typically greater than 5m for small lakes and 10m for larger ones) during summer, the water separates into layers of distinctly different density caused by differences in temperature, this process is called thermal stratification.

When the surface water cools again to approximately the same temperature as the lower water, the stratification is lost and the wind can turbulently mix the two water masses together because their densities are so similar. A similar process also may occur during the spring as colder surface waters warm to the temperature of bottom waters and the lake mixes. The lake mixing associated with a turnover often corresponds with changes in many other chemical parameters and this can affect biological communities, The National Advisory Committee to Water on the Web (2001).

For aquatic systems the maximum recommended increase in the natural temperature range of any inland or marine water is 2°C, Waterwatch Manual (1994). Because the Wetland Inventory has recorded the first readings for many wetlands, this natural temperature range is largely unknown. As a general guide, temperatures between zero and 20 degrees are acceptable limits for Mediterranean climates, while temperatures over 20 degrees can start impacting on aquatic fauna and fauna.

21.4.1 Analysis

The lowest water temperature recorded was 15 degrees, (Big White Salt Lagoon, Breakneck River) and the highest 28 degrees was recorded at Nepean Bay Lagoon. The most frequently recorded temperature was 26 degrees (6 sites). Generally there was an even spread of temperatures across all the wetland surveyed. Temperature readings are required over several time periods and during different seasons to attain a true indication of temperature regimes for wetlands on Kangaroo Island.



Note the algae growth over the water and the shallow water depth indicated by the gently sloping banks. The water temperature for this wetland was 28 degrees, the highest recorded in the Wetland Inventory.

Plate 21. Nepean Bay Lagoon

21.5 Dissolved oxygen

21.5.1 Background

Adequate concentrations of dissolved oxygen are necessary for the life of fish and other aquatic organisms. Dissolved oxygen levels (DO) are considered the most important and commonly employed measurement of water quality and indicator of a water body's ability to support aquatic life. Levels above 5 milligrams per liter (mg O₂/L) are considered optimal and most fish cannot survive for prolonged periods at levels below 3 mg O₂/L.

Seasonal changes affect dissolved oxygen concentrations. Warmer temperatures during summer speed up the rates of photosynthesis and decomposition. Pollution can also cause a decrease in dissolved oxygen concentrations.

The Australian and New Zealand Guidelines for Fresh and Marine Water Quality 2000 states that there is very little data on oxygen tolerances of Australian aquatic organisms. The 1992 ANZECC guidelines recommended that dissolved oxygen should not fall below 6 mg/L. The

National Science Foundation in America has developed the following guidelines for tolerance levels for aquatic invertebrates, these guidelines provide a means to analysis the survey results.

No production impairment 8 mgL
Moderate production impairment 5 mgL
Limit to avoid acute mortality 4 mgL

21.5.2 Analysis

Five sites recorded dissolved oxygen levels below 4 mgL (eg White Lagoon). Ten sites recorded dissolved oxygen levels between 5 mgL and 8 mgL (eg Grassdale Lagoon) and ten sites recorded dissolved oxygen levels of 8 mgL or more (eg Lashmar Lagoon). The highest recorded dissolved oxygen level (17 mg/L) was at Rush Lagoon, the lowest reading of 0.6 was recorded at Cygnet River Lagoon.



Rush Lagoon recorded the highest dissolved oxygen reading of 17 mg/L. This lagoon has high levels of aquatic vegetation contributing to the high oxygen levels.

Plate 22. Rush Lagoon.



Cygnet River Lagoon recorded the lowest dissolved oxygen reading of 0.6. It is interesting that a diversity of invertebrates were sampled from this lagoon.

Plate 23. Cygnet River Lagoon.

22.0 RAPID ASSESSMENT

The rapid assessment component of the survey provides a quick snap shot of the condition within different riparian habitats. This score is a subjective one given by the observer, it is an estimate of the percentage of intact native vegetation and the diversity and abundance of aquatic flora and fauna.

22.1 Aquatic Fauna

This is a subjective score according to the abundance and diversity of invertebrates, fish and birds located within the waterbody.

- Low (1) refers to no fish or birds with little or no invertebrate presence,
- Moderate (3) refers to some bird presence and good vertebrate abundance, fish usually are not present
- High (5) indicates good bird and invertebrate diversity and abundance with the presence of fish.




Figure 22-1. Aquatic Fauna Rapid Assessment

Twelve wetland sites recorded low aquatic fauna scores. This included wetlands such as Lashmar Lagoon, Milky's Lake, White Lagoon and Halls Road Swamp. Eleven wetlands scored as having moderate aquatic fauna (eg Murray Lagoon). Three wetlands recorded a high aquatic fauna presence (Breakneck River, Six-mile Lagoon and Rush Lagoon).

22.2 Aquatic flora

The rapid assessment for aquatic flora subjectively records the abundance and diversity of aquatic vegetation. The following scoring method has been developed:

- Low (1) indicates no or very little aquatic vegetation,
- Moderate (3) indicates some aquatic vegetation cover either in the form of floating or rooted vegetation
- High (5) indicates good diversity of aquatic vegetation with a range of rooted vegetation such as reeds and rushes and floating vegetation such as water ribbons.




Figure 22-2. Aquatic Flora Rapid Assessment

Fifteen sites recorded low aquatic vegetation (eg Salt Lagoon and Salt Lake) nine sites recorded moderate aquatic vegetation (eg White Lagoon) and two sites recorded high aquatic vegetation present (Murray Lagoon and Rush Lagoon).

22.3 Riparian Vegetation Rapid Assessment

Riparian native vegetation is assessed in three zones. These are the toe of bank (low water to high water), bank (high water mark to buffer) and buffer (top of the bank to the buffer which can extend between 10 metres and 100 metres). The vegetation association for each zone consists of the dominant or co-dominant overstorey species and understorey species. The vegetation association for each zone is scored on the basis of the level of disturbance and vegetation cover within each zone.

The vegetation association scoring system consists of:

- Degraded or no vegetation (1) with less than 30% vegetation cover, and a high level of disturbance
- Natural (3) between 30 – 75% vegetation cover, with little disturbance
- Intact (5) over 75% vegetation cover with little or no disturbance.

22.3.1 Analysis - Toe of bank




Figure 22-3. Rapid Assessment, Low – high water.

Fourteen sites recorded degraded banks at the high to low water mark (eg Discovery Lagoon, Lashmar lagoon, Moores Lake and Duck lagoon). Seven sites recorded a natural vegetation cover (eg White Lagoon, Salt Lake and Chapman River) and six sites recorded intact vegetation cover (eg Grassdale Lagoon, Six-mile Lagoon and Murray Lagoon).



Lashmar Lagoon recorded a degraded low to high water mark. The photograph illustrates this with no vegetation cover and signs of soil compaction and erosion from stock access.

Plate 24. Lashmar Lagoon.

23.3.2 Bank




Figure 22-4. Rapid Assessment, Bank

Nine sites recorded banks that are degraded (eg Birchmore Lagoon), 12 sites recorded banks with vegetation cover between 30 – 70% (eg Emanuel Lagoon), and six sites recorded banks with intact vegetation cover (eg Salt Lagoon).

23.3.3 Buffer




Figure 22-4. Rapid Assessment, Buffer

Eighteen sites recorded degraded buffers (eg Birchmore Lagoon), six sites recorded a natural vegetation cover (eg Murray Lagoon, Rush Lagoon) and three sites recorded intact buffers (Grassdale Lagoon, Breakneck River and Bay Lake 2). The wetlands with intact buffers are all located within NPWSA reserves.

23.0 WETLAND CONDITION SCORE

The wetland score reflects the previous rapid assessment scores for aquatic fauna, aquatic vegetation and riparian vegetation values. The combination of these values and the interpretation of other parameters recorded during the survey (such as land degradation and water chemistry) form the basis of the wetland condition score.

The wetland score consists of:

- Degraded (1) those sites that have a high level of disturbance and received low rapid assessment scores.
- Natural (3) those sites that have little disturbance, received moderate to high rapid assessment scores and that are sites usually located within National Parks and Wildlife reserves or managed on private lands for conservation purposes.
- Intact (5) those sites with no obvious sign of disturbance, scored very highly in the rapid assessment and are formally conserved within National Parks and Wildlife reserves.




Figure 23-1. Wetland score

Fourteen sites recorded a degraded wetland (eg Wangara Lagoon), nine sites recorded as having a natural wetland condition (eg Rush Lagoon) and three sites were recorded as being pristine (Grassdale Lagoon, Breakneck River and Bay Lake 2).



Grassdale Lagoon recorded a pristine wetland condition score. This lagoon has no major disturbances and scored highly in the rapid assessments. It is conserved within the NPWSA reserve system.

Plate 25. Grassdale Lagoon

24.0 Rapid assessment total score

This total score provides an indication of the environmental attributes of wetlands surveyed. These values are an interpretation of the biological, chemical and physical parameters. This score consist of a tally from all the rapid assessment scores, including the wetland condition scores, the scores consist of:

- 0 – 6 low wetland values
- 7 – 18 moderate wetland values
- 19 – 30 high wetland values

These scores provided a quick means of assessing the environmental attributes of wetlands based on the survey parameters. Those wetlands scoring high wetland scores should be more closely examined in terms of land ownership, placement within the landscape and the likely success of monitoring and protective measures.




Figure 24-1. Rapid assessment scores.

Six wetlands recorded low values (eg Lashmar Lagoon, Milkys Lake, Moores Lake), fifteen sites recorded moderate wetland values (eg Birchmore Lagoon, Emanuel Lagoon) and six sites recorded high wetland values and these are listed below.

Table 24-1. Wetlands with high rapid assessment scores.

Wetland name	Management Authority
Grassdale Lagoon	Department for Environment and Heritage
Breakneck River	Department for Environment and Heritage
Bay Lake 2	Department for Environment and Heritage
Six-mile Lagoon	Private
Murray Lagoon	Department for Environment and Heritage
Rush Lagoon	Private

25.0 ANZECC WETLAND CRITERIA

These criteria are used for determining nationally important wetlands in Australia, and forms the basis for the inclusion in the Directory for Important Wetlands. These criteria are those agreed to by the ANZECC Wetlands Network in 1994.

A wetland may be considered nationally important if it meets at least one of the following criteria:

1. *It is a good example of a wetland type occurring within a biogeographic region in Australia.*
2. *It is a wetland which plays an important ecological or hydrological role in the natural functioning of a major wetland system/complex.*
3. *It is a wetland which is important as the habitat for animal taxa at a vulnerable stage in their life cycles, or provides a refuge when adverse conditions such as drought prevail.*
4. *The wetland supports 1% or more of the national populations of any native plant or animal taxa.*
5. *The wetland supports native plant or animal taxa or communities which are considered endangered or vulnerable at the national level.*
6. *The wetland is of outstanding historical or cultural significance.*

Many of the sites in the Directory meet more than one of the criteria. Application of the criteria to individual wetland sites involves a degree of subjectivity. Aspects of a site's significance can be interpreted differently by different surveyors, and information gaps often exist which make it difficult to judge whether or not a site meets a particular criterion.

The Interim Biogeographic Regionalisation for Australia (IBRA) is used as the framework for applying Criterion 1, which identifies wetlands that are unique or representative within a biogeographic region in Australia.

25.1 Analysis

Sixteen sites recorded Criterion 1 (*good example of a wetland type occurring within a biogeographic region in Australia*). One site, Grassdale Lagoon, recorded Criteria 2 and two sites recorded Criteria 3, (Breakneck River and Grassdale Lagoon). Those wetlands with ANZECC Criteria are listed below.

Table 25-1. Wetlands with ANZECC Criteria.

Wetland Name	ANZECC Criteria
Salt Lagoon	1
Chapman River	1
Six-mile Lagoon 2	1
Grassdale Lagoon	1,2,3
Destree Swamp	1
White Lagoon	1
Rush Lagoon	1
Salt Lake 2	1
Bay Lake 2	1
Larrikin Lagoon	1
Lake Ada	1
Murray Lagoon	1
Lashmar Lagoon	1
Six mile Lagoon	1
Duck Lagoon	1
Breakneck River	1,3



Plate 26. Breakneck River



Plate 27. Grassdale Lagoon

Both Grassdale Lagoon and Breakneck River recorded multiple ANZECC criteria. Both aquatic systems are in good condition and have very good wetland values.

26.0 WETLAND TYPES

The definition of a wetland used in the survey is one adopted by the Ramsar Convention under Article 1.1.

“Wetlands are areas of marsh, fen, peatland or water, whether natural or artificial, permanent or temporary, with water that is static or flowing, fresh, brackish or salt, including areas of marine water the depth of which at low tide does not exceed six meters.”

Within this definition, the wetland classification system used in the Directory of Important Wetlands identifies 40 different wetland types in three categories: A-Marine and Coastal Zone wetlands, B-Inland wetlands, and C-Human-made wetlands. This wetland survey does not include Category A – Marine and Coastal Zone wetlands .

The system is based on that used by the Ramsar Convention in describing Wetlands of International Importance, but was modified slightly to suit the Australian situation.

A – Marine and Coastal Zone wetlands

1. Marine waters - permanent shallow waters less than six metres deep at low tide, includes sea bays straits
2. Subtidal aquatic beds, includes kelp beds, sea-grasses, tropical marine meadows
3. Coral reefs
4. Rocky marine shores, includes rocky offshore islands, sea cliffs
5. Sand, shingle or pebble beaches, includes sand bars, spits, sandy islets
6. Estuarine waters, permanent waters of estuaries and estuarine systems of deltas
7. Intertidal mud, sand or salt flats
8. Intertidal marshes, includes salt marshes, salt meadows, saltings, raised salt marshes, tidal brackish and freshwater marshes
9. Intertidal forested wetlands, includes mangrove swamps, nipa swamps, tidal freshwater swamps forests
10. Brackish to saline lagoons and marshes with one or more relatively narrow connections with the sea
11. Freshwater lagoons and marshes in the coastal zone
12. Non tidal freshwater forested wetlands

B – Inland wetlands

1. Permanent rivers and streams includes waterfalls
2. Seasonal and irregular rivers and streams
3. Inland deltas(permanent)
4. Riverine floodplains, includes river flats, flooded river basins, seasonally flooded grassland, savanna and palm savanna
5. Permanent freshwater lakes (8 ha) includes large oxbow lakes
6. Seasonal/intermittent freshwater lakes (>8 ha) floodplain lakes
7. Permanent saline /brackish lakes
8. Seasonal/intermittent saline lakes
9. Permanent freshwater ponds (<8 ha) marshes and swamps on inorganic soils, with emergent vegetation waterlogged for at least most of the growing season
10. Seasonal/intermittent freshwater ponds and marshes on inorganic soils includes sloughs, potholes, seasonally flooded meadows, sedge marshes
11. Permanent saline/brackish marshes
12. Seasonal saline marshes
13. Shrub swamps, shrub dominated freshwater marsh, shrub carr, alder thicket on inorganic soil
14. Freshwater swamp forest, seasonally flooded forest, wooded swamps, on inorganic soils
15. Peatlands, forest, shrubs or open bogs
16. Alpine and tundra wetlands: includes alpine meadows, tundra pools, temporary waters from snow melt
17. Freshwater springs, oasis and rock pools
18. Geothermal wetlands
19. Inland, subterranean karst wetlands

C- Human-made wetlands

1. Water storage areas; reservoirs, barrages, hydro-electric dams, impoundment's (generally over 8 ha).
2. Ponds; includes farm ponds, stock ponds, small tanks; (generally below 8 ha).
3. Aquaculture ponds; fish ponds shrimp ponds
4. Salt exploitation, salt pans, salines
5. Excavations; gravel pits; borrow pits, mining pools.
6. Wastewater treatment areas; sewage farms, settling ponds, oxidation basins.
7. Irrigated land; includes irrigation channels and rice fields, canals, ditches
8. Seasonally flooded arable land, farm land
9. Canals

26.1 Analysis




Figure 26-1. Wetland Types in Kangaroo Island

Permanent saline/brackish lakes are recorded 11 times, including Salt Lake and Big White Salt Lagoon. Seasonal/intermittent saline lakes (B8) are recorded nine times and include wetlands such as Nepean Bay Lagoon.



Plate 28. Bay Lake

Bay Lake (wetland type B10) located within Flinders Chase National Park is a good example of a seasonal freshwater pond.



Plate 29. Emanuel Lagoon

Emanuel Lagoon is an example of wetland type B8 (Seasonal/intermittent saline lakes).

SECTION THREE WETLAND MONITORING

27.0 INTRODUCTION

There are three important steps in developing monitoring protocols that can be applied effectively within the South Australian wetland management context.

These consist of:

1. Identification of wetland values (collating and collecting existing information and undertaking baseline surveys)
2. Identification of wetland threats (from the analysis of wetland values)
3. Development of monitoring indicators that highlight early changes in ecological Character.

“Monitoring essentially means the keeping of a continual record of certain parameters, advising whether they are being maintained within prescribed limits and warning if undesirable changes occur” (Hart 1980).

Finlayson and Eliot (2001) expand on this definition by emphasising that the information derived from monitoring provides a platform on which management actions are both based and judged.

Finlayson and Mitchell (1999) identified the following five major reasons for monitoring wetlands:

- To characterise variations in responses of wetlands to natural variability in the environment;
- To collect baseline data on wetlands as part of inventory processes;
- To record ecological changes that may be occurring as a result of specific natural or anthropogenic events;
- To measure progress toward set objectives of a management program; and
- To audit the performance of management agencies and land users.

27.1 Ecological change

The Ramsar Convention for Internationally Important Wetlands states that in order to record ecological change, the ecological character needs to be defined. Once defined it is possible to monitor for changes in ecological character.

Ecological character is the sum of the biological, physical, and chemical components of the wetland ecosystem, and their interactions, which maintain the wetland and its products, functions, and attributes. The process of completing a wetland inventory assists in establishing the ecological character for wetlands surveyed.

The change in ecological character is the impairment or imbalance in any biological, physical, or chemical components of the wetland ecosystem, or in their interactions, which maintain the wetland and its products, functions and attributes. (www.ramsar.org/key_guide-list-e.html)

The Ramsar Convention provides detailed guidelines for monitoring the ecological character of wetlands. These include guidelines for initiating and conducting a risk assessment framework. The Ramsar risk assessment framework consists of six steps, these are:

Step 1 - Identification of the problem. This is the process of identifying the nature of the problem and developing a plan for the remainder of the risk assessment based on this information. It defines the objectives and scope of, and provides the foundation for, the risk assessment. In the case of a chemical impact, it would include obtaining and integrating information on the characteristics (for example, properties, known toxicity) and source of the chemical, what is likely to be affected, and how is it likely to be affected, and importantly, what is to be protected.

Step 2 - Identification of the adverse effects. This step evaluates the likely extent of adverse change or impact on the wetland. Such data should preferably be derived from field studies, as field data are more appropriate for assessments of multiple impacts, such as occur on many wetlands. Depending on the extent of adverse change and available resources, such studies can range from quantitative field experiments to qualitative observational studies. For chemical impacts, on-site ecotoxicological bioassays constitute appropriate approaches, whereas for changes caused by weeds or feral animals, on-site observation and mapping may be all that is required.

Step 3 - Identification of the extent of the problem. This step estimates the likely extent of the problem on the wetland of concern by using information gathered about its behaviour and extent of occurrence elsewhere. In the case of a chemical impact, this includes information on processes such as transport, dilution, partitioning, persistence, degradation, and transformation, in addition to general chemical properties and data on rates of chemical input into the environment. In the case of an invasive weed, it might include detailed information on its entry into an ecosystem, rate of spread and habitat preferences. While field surveys most likely represent the ideal approach, use of historical records, simulation modeling, and field and/or laboratory experimental studies all represent alternative or complementary methods of characterising the extent of the problem.

Step 4 - Identification of the risk. This involves integration of the results from the assessment of the likely effects with those from the assessment of the likely extent of the problem, in order to estimate the likely level of adverse ecological change on the wetland. A range of techniques exist for estimating risks, often depending on the type and quality of the likely effects and their extent. A potentially useful technique for characterising risks in wetlands is via a GIS-based framework, whereby the results of the various assessments are overlaid onto a map of the region of interest in order to link effects to impact. In addition to estimating risks, such an approach would also serve to focus future assessments and/or monitoring on identified problem areas.

Step 5 - Risk management and reduction. This is the final decision-making process and uses the information obtained from the assessment processes described above, and it attempts to minimize the risks without compromising other societal, community or environmental values. In the context of the Ramsar Convention, risk management must also consider the concept of *wise use* and the potential effects of management decisions on this. The result of the risk assessment is not the only factor that risk management considers; it also takes into account political, social, economic, and engineering/ technical factors, and the respective benefits and limitations of each risk-reducing action. It is a multidisciplinary task requiring communication between site managers and experts in relevant disciplines.

Step 6 - Monitoring. Monitoring is the last step in the risk assessment process and should be undertaken to verify the effectiveness of the risk management decisions. It should incorporate components that function as a reliable early warning system, detecting the failure or poor performance of risk management decisions prior to serious environmental harm occurring. The risk assessment will be of little value if effective monitoring is not undertaken. The choice of endpoints to measure in the monitoring process is critical. Further, a GIS-based approach will most likely be a useful technique for wetland risk assessment, as it incorporates a spatial dimension that is useful for monitoring adverse impacts on wetlands.

28.0 MONITORING PROTOCOLS AND INDICATORS

The underlying concept of indicators is that effects can be detected before actual environmental impacts occur. While such a 'early warning' may not necessarily provide firm evidence of larger scale environmental degradation, it provides an opportunity to determine whether intervention or further investigation is warranted.

Early warning indicators can be defined as:

"The measurable biological, physical or chemical responses to a particular stress, preceding the occurrence of potentially significant adverse effects on the system of interest".
http://ramsar.org/key_guide_risk_e.htm

Typically, indicators are ecological communities or assemblages of organisms, habitat or keystone-species. Indicators are usually closely linked to ecosystem-level effects. In selecting an indicator it is important that the ecological character of a wetland which includes the biological, chemical and physical components of the ecosystem. Therefore, it may be useful to select early warning indicators according to which of the above three components are considered more susceptible to change within the wetland, Finlayson and Spiers (1999).

The concepts of early warning and ecological relevance can conflict. As an example, biomarker responses can offer exceptional early warning of potential adverse effects. However, there exists very little evidence that observed responses result, or culminate in adverse effects at an individual level, let alone the population, community or ecosystem level Finlayson and Spiers (1999). If the primary assessment objective is that of early detection, then it is likely that it will be at the expense of ecological relevance (Finlayson and Spiers, 1999).

van Dam *et al* (*in press*) in Finlayson and Spiers (1999), suggests early warning indicators should be:

1. **anticipatory**: it should occur at levels of organisation, either biological or physical, that provide an indication of degradation, or some form of adverse effect, before serious environmental harm has occurred;
2. **sensitive**: in detecting potential significant impacts prior to them occurring, an early warning indicator should be sensitive to low levels, or early stages of the problem;
3. **diagnostic**: it should be sufficiently specific to a problem to increase confidence in identifying the cause of an effect;
4. **broadly applicable**: it should predict potential impacts from a broad range of problems;
5. **correlated to actual environmental effects/ecological relevance**: an understanding that continued exposure to the problem, and hence continued manifestation of the response, would usually or often lead to significant environmental (ecosystem-level) adverse effects;
6. **timely and cost-effective**: it should provide information quickly enough to initiate effective management action prior to significant environmental impacts occurring, and be inexpensive to measure while providing the maximum amount of information per unit effort;

7. **regionally or nationally relevant:** it should be relevant to the ecosystem being assessed;

8. **socially relevant:** it should be of obvious value to, and observable by stakeholders, or predictive of a measure that is socially relevant;

9. **easy to measure:** it should be able to be measured using a standard procedure with known reliability and low measurement error;

10. **constant in space and time:** it should be capable of detecting small change and of clearly distinguishing that a response is caused by some anthropogenic source, not by natural factors as part of the natural background (that is, high signal to noise ratio);

11. **nondestructive:** measurement of the indicator should be nondestructive to the ecosystem being assessed.

Finlayson and Spiers (1999) discuss that the importance of the above attributes cannot be over-emphasized, since any assessment of actual or potential change in ecological character will only be as effective as the indicators chosen to assess it. However, an early warning indicator possessing all the ideal attributes cannot exist, as in many cases some of them will conflict, or will simply not be achievable.

28.1 Early warning indicators

Finlayson and Spiers (1999) outline a number of early warning indicators, these are placed into three broad categories:

1. rapid response toxicity tests;
2. field early warning tests; and
3. rapid assessments.

Each of the techniques may meet different objectives in water quality assessment programs. Although the majority of early warning indicators are of a biological nature, physico-chemical indicators do exist and often form the initial phase of assessing water quality (Finlayson and Spiers, 1999).

28.2 Toxicity tests

These represent rapid and sensitive responses to one or more chemicals. They provide an indication that there may be a risk of adverse effects occurring at higher levels of biological organization (for example, communities and ecosystems). Laboratory toxicity tests are of particular use for a chemical or chemicals yet to be released into the aquatic environment (for example, a new pesticide or a pre-release waste water). They provide a basis upon which to make decisions about safe concentrations or dilution/release rates, thereby eliminating, or at least minimizing, adverse impacts on the aquatic environment. However, there are major differences in the ecological relevance of responses that can be measured, Finlayson and Spiers (1999).

28.3 Field early warning indicators

These comprise of a range of techniques that are grouped together because they are used to measure responses or patterns in the field and thus provide a more realistic indication of effects in the environment. In contrast to rapid response toxicity tests, early warning field tests predict and/or assess the effects of existing chemicals. Some of the techniques can also be applied to biological and physical problems (Finlayson and Spiers, 1999).

28.4 Rapid Assessments

Rapid assessment is essentially the method used for this wetland inventory, and has the appeal of enabling ecologically-relevant information to be gathered over wide geographical areas in a standardised fashion and at relatively low costs. The trade-off is that rapid assessment methods are usually relatively 'coarse' and hence are not designed to detect subtle impacts.

Finlayson and Spiers (1999) point out that the attributes of rapid assessment include:

1. measured response is widely regarded as adequately reflecting the ecological condition or integrity of a site, catchment or region (that is, ecosystem surrogate);
2. approaches to sampling and data analysis are highly standardised;
3. response is measured rapidly, cheaply and with rapid turnaround of results;
4. results are readily understood by non-specialists; and
5. response has some diagnostic value.

The most powerful impact assessment programs will generally be those that include two types of indicator, namely those associated with early warning of change and those closely associated with ecosystem-level effects. The ecosystem-level indicator might include ecologically important populations (keystone species). With both types of indicators used in a monitoring program, information provided by ecosystem-level indicators may then be used to assess the ecological importance of any change observed in an early detection indicator.

29.0 RECOMMENDED INDICATORS FOR MONITORING SURVEYED WETLANDS

Monitoring **water chemistry** is recommended as an **early warning indicator** and monitoring **invertebrate composition** as an **ecosystem level indicator**.

Before these indicators can be used effectively in a monitoring program, more research and surveys are required. The wetland surveys undertaken on Kangaroo Island recorded information rapidly at one location and at one time of the year. An understanding of the natural levels of water chemistry and invertebrate composition for selected wetlands is needed before effective monitoring can occur.

29.1 Early warning indicators – Water chemistry

The collection of water chemistry parameters is suggested as the means for monitoring for early change in wetlands. Parameters essential to collect include conductivity, dissolved oxygen, temperature, turbidity and pH. Conductivity will probably play an important indicator providing the first early warning change within wetlands.

29.2 Ecosystem based indicator

Invertebrates are recommended as indicators for ecosystem monitoring. Yen & Butcher (1997) outline the important features of invertebrates which make them useful as indicators. These include;

1. Invertebrates are ecologically and functionally important.
2. Invertebrates are diverse, providing a good range from which to choose suitable taxa.
3. Many species are habitat specific.
4. Many species are abundant and are relatively simple to collect.

Indicator taxa should be selected on a regional basis and that are sensitive to environmental change. Indicator species should also be selected that have been identified as keystone species, these species play critical ecological roles, and their loss may result (directly or indirectly) in the disappearance of several other species, Soule and Kohm, 1989); Lawton, 1991 in Yen and Butcher, 1997).

Research is required towards developing these indicators. Yen and Butcher (1997) suggest research in the following areas:

- The link between invertebrate biodiversity and ecosystem functioning.
- The value of using guilds or functional groups to monitor ecosystem processes.
- Identifying keystone taxa.
- The existence of redundant species.

The interaction between invertebrate composition and water chemistry also requires more research. Without an understanding of the dynamics between these two parameters, interpretation of ecological change will be hindered.

An example of how the invertebrate monitoring program can be applied to detect change in ecological character is to study the trophic dynamics or functional groups. The presence of both invertebrate grazers and predators indicates functional invertebrate dynamics. The loss of either group quickly indicates change in the ecological character of the wetland.

The development of keystone indicator species requires further research, However, as a starting point there are several families that can be used in starting to develop these indicators. Representative grazers and predators are important to include as indicators due to their importance in the trophic chain.

Keystone species to monitor include:

Grazers: zooplankton, including Cladocera, Copepods, Calanoida and Cyclopoida.

Predators: dragonfly larvae (Odonata) and the families Dytiscidae and Coleoptera.

29.3 Other considerations

Monitoring is also required for many other parameters. Programs need to include whole of catchment parameters that cause change in the ecological character of wetlands. Monitoring of threats to wetland such as land clearance and management, drainage, land ownership changes and land use changes all need to be documented and included in the monitoring program. Ideally an action plan to alleviate and manage threats should be developed for nominated wetlands in conjunction with a monitoring program.

30.0 RECOMMENDED WETLANDS TO MONITOR

Those wetlands that received a high wetland score and fulfilled the ANZECC criteria for an important wetland are priority wetlands to monitor. The second criteria used to identify priority wetlands the management authority. Those wetlands that are under public ownership are targeted as preferential sites to monitor unless significant wetland values are identified on private lands. The reasoning behind this is that wetlands within the community land classification or under State government ownership may be easier to access and manage a monitoring program. At this early stage of developing a monitoring program the nominated sites will assist in refining techniques and develop monitoring protocols.

Once monitoring techniques have been developed and tested, the opportunity to expand monitoring to other wetlands with high conservation value on private property is feasible. Other protection measures can be developed for wetlands on private property before monitoring occurs. Actions such as buffering wetlands with revegetation, fencing, removing rubbish and the development of wise use of wetland programs for landowners are currently required to manage wetland threats. The following wetlands that are recommended for monitoring cover a broad range of wetland types, display different water regimes, vegetation structure and geographical locations.

30.1 Recommended priority wetlands to monitor

Table 30-1. Recommended wetlands to monitor

Wetland site number	Wetland name	Wetland features	Management Authority
S5071	Grassdale Lagoon	Intact freshwater wetland system	Department for Environment and Heritage
S5056	Six mile Lagoon	Small inclosed freshwater wetland, with high diversity of invertebrates	Private
S5018	Rush Lagoon	Large saline lake with high aquatic productivity	Private

31.0 RECORDING MONITORING PARAMETERS

The wetland inventory template (refer to [Section One](#) and [Appendix 1](#)) should be used for the collection of baseline data. Once collected, analysis of the results is required to determine the values of the wetland, threats and possible key indicator groups. Routine collection of information is required in order to establish which monitoring indicators to use (early warning and ecosystem indicators). After a given time period (to be determined for each site comparison of the data is required to identify if any ecological change has occurred. Defining acceptable limits of ecological change will be difficult and may take some time until the wetland processes are understood. The suggested wetlands to monitor will be of value in developing reference sites for other wetlands in the region. Each of the three wetlands suggested to monitor represent different wetland types, within different landforms and under a range of landownerships.

BIBLIOGRAPHY

- A.N.Z.E.C.C (1999). *Draft Overview of the National Water Quality Management Strategy.*, Australian and New Zealand Guidelines for Freshwater and Marine Water Quality.
- Arthington, M. K. (1997). *Review of impacts of displaced/introduced fauna associated with inland waters.* Canberra, Department for Environment: 69.
- Barritt, M. (1982). *A rapid appraisal survey of some natural areas on kangaroo Island, South Australia.* Adelaide, Nature Conservation Society.
- Bek, P. A.. Robinson, G. (1991). *Sweet Water or Bitter Legacy-Water Quality in New South Wales,* Water Quality Unit, Department of Water Resources, Sydney.
- Biological Database of SA; *Reserves and plant populations;* Flora. Source for status; SA = NPWSA Act 1972, Updated Schedule 1999, REG = Lang and Kraehenbuehl, 2001.
- Brock, M.(1999). *Australian freshwater ecology: processes and management,* Gleneagles Publishing.
- Brown, W.(1979). *An inventory of research and available information on wetlands in Western Australia,* Department of Conservation and Environment, WA.
- Butcher, R. (1999). *Assessing biodiversity in temporary and permanent wetlands,* Pp 50-53 in *The Other 99%. The Conservation and Biodiversity of Invertebrates,* ed by W. Ponder and D. Lunney, 1999. Transactions of the Royal Zoological Society of New South Wales, Mosman.
- Butcher, Y.(1997). *An overview of the conservation of non-marine invertebrates in Australia.* Canberra, Environment Australia.
- Cadwallader, B (1983). *A guide to the freshwater fish of Victoria,* FD Atkinson Government Printer.
- Curruthers, S. and Hille, B. (1997) *South East Wetlands GIS Database,* Geographic Analysis and Research Unit, Information and Data Analysis Branch, Planning Division, Department of Housing and Urban Development.
- Davis, M.(1998). *Wetlands for the future,* Gleneagles Publishing, SA.
- Defaye, D.(1995). *Copepoda, Introduction to the Copepoda,* SPB Academic Publishing.
- DEH (2001) *Provisional listing of threatened ecosystems of the Agricultural Regions for South Australia.*
- Department of Conservation and Environment (1980). *Guidelines for the conservation and management of wetlands in Western Australia.*, Department of Conservation and Environment, Western Australia.
- Department of Housing and Urban Development, (1997). *Guide to a native vegetation survey using the biological survey of South Australia methodology, Section 3,* Geographic Analysis and Research Unit.
- Dowall., R.(1980). *Freshwater fishes of South-Eastern Australia,* A.H. & A.W. Reed PTY LTD.
- Dussart, B. H.(2001). *Introduction to the Copepoda,* Backhuys Publishers.
- Environment Protection and Biodiversity Conservation Act, 1999
- Environmental Protection Authority (1993). *A guide to wetland management in the Swan coastal plain area.*, Environmental Protection Authority, Perth, Western Australia.
- Fairweather, P.G. & Napier (1998). *Environmental indicators for national state of the environment reporting - inland waters.* Environment Australia.
- Farinha, N.(1996). *Mediterranean Wetland Inventory: A Reference Manual.*, MedWet/Instituto da Coservacao da Natureza/Wetlands International Publication, Volume 1, Lisbon and Slimbridge, UK.
- Finlayson, C.M. and Spiers, A.G. (1999). *Techniques for enhanced wetland inventory and monitoring.* Supervising Scientist, Canberra.
- Finlayson, C.M. and Eliot, I. (2001). "Ecological Assessment and monitoring of Coastal Wetlands in Australia's Wet Dry Tropics: A Paradigm for elsewhere?" *Coastal Management* 29: 105-115.
- Finlayson, C. M. and Mitchell, D.S. (1999). "Australian wetlands: The monitoring challenge." *Wetlands ecology and management* 7: 105-112.
- Geneva, W. H. O (1996). *Guidelines for drinking-water quality,* 2nd ed. Vol. 2. Health criteria and other supporting information pp. 370-373.
- Greenway, M. (1997) *Wetlands in a Dry Land: Understanding for Management,* Albury, New South Wales, 29-30. Environment Australia.
- Hart, B. T. (1980). *Environmental surveys. An Ecological Basis for Water Resource Management,* Australian National University Press.

- Heard and Channon (1997) Guide to a native vegetation survey using the biological survey of South Australia methodology, Section 3. Geographic Analysis and Research Unit, Department of Housing and Urban Development.
- Jackson, J.(1997). *State of Habitat Availability and Quality in Inland Waters*. Australia: State of the Environment Technical Paper Series (Inland Waters), Environment Australia.
- Jensen, A.(1996). *Wetlands Atlas of the South Australian Murray Valley*, Department for Environment and Heritage.
- Liston and Maher, (1997). *Water quality for maintenance of aquatic ecosystems: Appropriate indicators and analysis*. Australia: State of the Environment Technical Paper Series. (Inland waters)., Environment Australia.
- Lloyd, B.(1986). *Wetlands & Water Resources of South Australia*. Appendix 1. Part 3: SouthAustralian Gulf Drainage Division., Conservation Projects Branch. Department of Environment & Planning.
- Ludbrock, N.H. (1980) *A guide to the geology and mineral resources of South Australia*. Department of Mines and Energy South Australia. Government Printer SA.
- Maher, W. & Liston, P. (1997). *Water quality for maintenance of aquatic ecosystems: Appropriate indicators and analysis*. Australia: State of the Environment Technical Paper Series. (Inland waters). Environment Australia.
- McComb, P.(1988). *The Conservation of Australian Wetlands*, Surrey Beatty & Sons Pty Limited, NSW.
- Morelli, J. and DeJong, M.(1996). *A Directory of Important Wetlands in South Australia*. Adelaide, South Australian Department of Environment and Natural Resources.
- Neagle, N. (2002) *National Land and Water Audit, Conservation Strategy Case Study, Kangaroo Island Subregion, SA*. Department for Environment and Heritage SA.
- Nogrady, T.(1993). *Rotifera, guides to identification of the Micro Invertebrates of the Continental Waters of the World.*, SPB Academic Publishing.
- NLWRA (National Land and Water Resources Audit) 2001, *Australian Dryland Salinity Assessment 2000. Extent, impacts, processes, monitoring and management options*, National Land and Water Resources Audit, Australia.
- Paijmans, K.(1978). *A Reconnaissance of four wetland pilot study areas.*, CSIRO Division of Land Use Research, Canberra.
- Pressey, R. L.(1987). *A Survey of wetlands of the lower Macleay Floodplain*, NSW, NSW National Parks and Wildlife.
- Raines, J.(1993). *Nomination of wetlands of outstanding ornithological importance for the Register of the National Estate*. Royal Australasian Ornithologist Union Database, Western Australia.
- Ramsar (*Ramsar Resolution VII.10 and Annex on Wetland Risk Assessment Framework*). Ramsar.
- Rivier, I. K.(1998). *The predatory Cladocera (Onychopoda: Podonidae, Polyphemidar, Cercopagidae) and Leptodorida of the world.*, Backhuys Publishers.
- Robinson, P.(1991). *Sweet Water or Bitter Legacy-Water Quality in New South Wales*, Water Quality Unit, Department of Water Resources, Sydney.
- Rolls, D. (1987). *A baseline biological monitoring programme for the urban wetlands of the Swan Coastal Plain*, WA. Perth, Environmental Protection Authority.
- Romanowski, Nick. (1998). *Aquatic and wetland plants*.
- Sainty, G. R. (1981). *Waterplants of New South Wales*, Water Resources Commission NSW.
- Sainty, G. R.(1994). *Waterplants in Australia*, South China Printing Co.
- Sheldon, F. and Walker, K.(2001). *"Propagules in dry wetland sediments as indicators of ecological health: effects of salinity."* Regulated rivers:Research and Management 17: 191-197.
- Smet, W.(1997). *Rotifera, Volume 5: The Dicranophoridae (Monogononta)*, SPB Academic Publishing.
- Smirnov (1992). *The Macrothricidae of the world*, SPB Academic Publishing.
- Sonneman, A.(1988). *Freshwater Algae in Australia*, Sainty and Associates.
- Stanton, J. P. (1975). *A Preliminary Assessment of Wetlands in Queensland*, CSIRO Division of Land Use Research, Canberra.
- Storey, A.W. Lane, J.A.K. Davies, P.M. (1997). *Monitoring the ecological character of Australia's wetlands of international importance (RAMSAR Convention)*. Western Australian Department of Conservation and Land Management and Biodiversity Group of Environment Australia.
- Streever, W.(1996). *Preliminary example of a sampling design assessment method for biomonitoring studies that rely on ordination*. INTECOL'S V International Wetlands Conference., University of Western Australia, Gleneagles Publishing.
- Thackway, R. and Cresswell, I. (1995). *An Interim Biogeographic Regionalisation for Australia: a framework for setting priorities*

in the National Reserves System Cooperative Program Version 4, Australian Nature Conservation Agency, Canberra.

The National Advisory Committee to Water on the Web (2001) <http://wow.nrri.umn.edu>

Thompson, M. B.(1986). *River Murray Wetlands:their characteristics, significance and management.*, Nature Conservation Society of South Australia and the Department of Environment and Planning, Adelaide.

Walker S.J. and Goonan P.M.(2000) *Frog Census 2000*, Environmental Protection Agency, DEH. South Australia.

Walker, G. van der Wel, B. & Leaney, R. Jolly, I.(2000). *Assessing the impacts of dryland salinity on South Australia's water resources*. South Australia, CSIRO Land and Water: 60.

Waterwatch Manual (1996). *Waterwatch Manual for South Australia*, South Australian Government Printer.

Weatherley, A. H.(1967). *Australian inland waters and their fauna*, Australian National University Press, Canberra.

Williams, W. D.(1980). *Australian Freshwater Life*, Globe Press, Victoria.

Williams, W. D.(1998). *Wetlands in a dry land: understanding for management.*, Environment Australia, Biodiversity Group, Canberra.

Williams, W.D. (2000). "Biodiversity in temporary wetlands of dryland regions." *Limnology* 27: 141-144.

Williams, W.D.(2001). *Salinisation:unplumbed salt in a parched landscape*.

Willoughby, N., Oppermann, A., and Inns, R.W. (2001) Biodiversity Plan for Kangaroo Island, South Australia. Department for Environment and Heritage, South Australia.

Willoughby, N. (1999). Preliminary to the Biodiversity Plan for Kangaroo Island. Department for Environment, Heritage and Aboriginal Affairs.

Web Sites referenced

http://www.ea.gov.au/parks/nrs/ibraimcr/ibra_95/index.html
<http://www.nmnh.si.edu/botany/projects/algae/Imag-Chl.htm>
http://www.who.int/water_sanitation_health/GDWQ/Chemicals/turbidity.htm
<http://www.epa.nsw.gov.au/soe/97/ch3/11.htm>
<http://wow.nrri.umn.edu/wow/under/parameters/turbidity.html>
www.ramsar.org/key_guide-list-e.html
http://ramsar.org/key_guide_risk_e.htm

Other useful web sites

Water Watch
<http://www.sa.waterwatch.org.au/>

Environment Australia
<http://www.ea.gov.au/water/wetlands/>

Wetland Care Australia
<http://www.wetlandcare.com.au/>

Environmental Protection Agency
<http://www.environment.sa.gov.au>

AusRivas
<http://ausrivas.canberra.edu.au/>

Catchment Water Management Boards
<http://www.catchments.net/>

Invertebrates
<http://www.umesci.maine.edu/ams/inverts.htm>

Appendix 1 Wetland Inventory descriptions

Wetland Reference Number: The wetland reference number is critical for clearly defining wetland locations within regions of the State. The wetland reference number acts as the primary key for records for each wetland, this enables data transfer between databases and GIS tables.

Wetland Name: The name of the site.

Ramsar Site: If the wetland is listed as a Ramsar site.

Description of Site: A general description of the location features, closest roads or landmarks.

Current Land use

On site: Current human use of the designated wetland area.

Surrounding area: Human use on land adjacent to the wetlands, and more broadly in the surrounding catchment.

Tenure/ownership

On site: Details of land ownership of the wetland site if possible.

Surrounding: Details of the tenure type that is dominant in the surrounding areas if possible.

Jurisdiction and management authority: The name of the body or bodies responsible for management of the wetland.

Complied by: Name of person undertaking data collection.

Organisation: Organisation that is managing the data collection

Date/time: Date and time of data collection.

Region: Geographical region (eg Eyre Peninsula, Yorke Peninsula and Kangaroo Island)

Environmental Regions: This refers to broad environmental regions within South Australia as described by Laut P (1977), or regions described within the Regional Biodiversity Plans developed by the Department for Environment and Heritage (2000 – 2001).

Key Bio diversity Areas: The Department for Environment and Heritage has developed key biodiversity areas within the Regional Biodiversity Plans for South Australia. The key biodiversity areas have been developed by analysis of biological assets of the area and information from community groups. There are six key biodiversity areas identified within the three wetland study regions; these are: coastal wetland habitat areas, fragmented habitat areas, grassland habitat, large remnant area, ridgeline systems, threatened habitat area.

GPS reading: Taken in Latitude and Longitude or easting and northing.

Map Sheet and Reference: Record of map sheet name and number with grid reference if required.

Approximate Area: usually recorded after or before survey by GIS query.

Elevation: in meters above sea level. Preferably recorded from GPS and compared with map sheet elevation values.

Landform Element: Landform element definitions have been adapted from the "Guide to a native vegetation survey using the biological survey of South Australia methodology, Section 3". (1997) Geographic Analysis and Research Unit, Department of Housing and Urban Development.

Sandy Plain: Large, very gently inclined or level element composed of fine grains of weathered rocks of quartz.

Limestone plain: Large, very gently inclined or level element of hard almost horizontally bedded limestone, which contains at least 80% of the carbonates of calcium or magnesium.

Rock outcrop: Any exposed area of rock that is inferred to be continuous with underlying bedrock on a large, very gently inclined or level element.

Drainage depression: Level to gently inclined, long, narrow, shallow open depression with smoothly concave cross-section, rising to moderately inclined side slopes, eroded or aggraded by sheet wash.

Open depression: Landform element that extends at the same elevation, or lower, beyond the locality where it is observed.

Closed depression: Landform element that stands below all points in the adjacent terrain.

Flat: Planar landform element that is neither a crest nor a depression and is level or very gently inclined (<3% slope).

Swale: Linear, level-floored open or closed depression excavated by wind, or left relict between ridges built up by wind or waves, or built up to a lesser height than them. Or long, curved open or closed depression left relict between scrolls built up by channelled stream flow.

Lagoon: Closed depression filled with water that is typically salt or brackish, bounded at least in part by forms aggraded or built up by waves or reef building organisms.

Hill Crest: Very gently inclined to steep crest, smoothly convex, eroded mainly by creep and sheet wash. A typical element of mountains, hills, low hills and rises.

Hill slope: Gently inclined to precipitous slope, commonly simple and maximal, eroded by sheet wash, creep, or water-aided mass movement.

Gully: Open depression with short, precipitous walls and moderately inclined to very gently inclined floor or small stream channel, eroded by channelled stream flow and consequent collapse and water-aided mass movement.

Stream channel: Linear, generally sinuous open depression, in parts eroded, excavated, built up and aggraded by channelled stream flow. This element comprises stream bed and banks.

Flood out: Flat inclined radially away from a point on the margin or at the end of a stream channel, aggraded by over bank stream flow, or by channelled stream flow associated

with channels developed within the over-bank flow part of a covered plain landform pattern.

Fan-alluvial: Large gently inclined to level element with radial slope lines inclined away from a point, resulting from aggradation, or occasionally from erosion, by channelled, often braided, stream flow, or possibly by sheet flow.

Estuary: Stream channel close to its junction with a sea or lake, where the action of channelled stream flow is modified by tide and waves. The width typically increases downstream.

Lake: Large water-filled closed depression.

Salt lake: Lake, which contains a concentration of mineral salts, (predominantly sodium chloride in solution as well as magnesium and calcium sulphate).

Swamp: Almost level closed, or almost closed depression with a seasonal or permanent water table at or above the surface, commonly aggraded by over-bank stream flow and sometimes biological (peat) accumulation.

Perched swamp: A tract of land that is permanently saturated with moisture and is positioned on an elevated landform.

Playa/pan: Large, shallow, level-floored closed depression, intermittently water-filled, but mainly dry due to evaporation, bounded as a rule by flats aggraded by sheet flow and channelled stream flow.

Geological formation name: Name of geological formation defined from the physical geography data sets available from <http://www.atlas.sa.gov.au/>

Soil types: Soil type definitions have been adapted from the "Guide to a native vegetation survey using the biological survey of South Australia methodology, Section 3". (1997) Geographic Analysis and Research Unit, Department of Housing and Urban Development. To determine soil texture take a small handful of soil below the crust and add water working the soil into a an elongated ball (bolus) until it just fails to stick to the fingers. The behaviour of the bolus and of the ribbon produced by shearing (pressing out) between the thumb and forefinger characterise the texture. The behaviour and feel, smoothness or graininess, during bolus formation are also indicative of its texture.

Sand: Coherence nil to very slight, cannot be moulded. Sand grains of medium size, single sand grains adhere to fingers. Commonly less than 5% clay content.

Loam: Bolus coherent and rather spongy, smooth feel when manipulated but with no obvious sandiness. May be somewhat greasy to the touch if organic matter in present. Will form a ribbon of about 25mm.

Clay Loam, Sandy: Coherent plastic bolus with medium size sand grains visible within a finer matrix, forms a ribbon between 40-50mm. Clay contents between 30% – 35%.

Medium clay: Smooth plastic bolus, can be moulded into rods without fracture. Will form a ribbon of 75 mm or more. Clay contents between 45% - 55%.

Medium heavy clay: Same properties as medium clay but with a higher clay content of 50% or more.

Loamy sand: Slight coherence, sand grains of medium size, can be sheared between thumb and forefinger to give minimal ribbon of about 5 mm. Approximate clay content of about 5%.

Silty loam: Coherent bolus, smooth, often silky when manipulated, will form a ribbon of about 25mm. Clay content between 25% with 25% or more of silt.

Silty clay loam: Coherent smooth bolus, plastic and often silky to the touch, will form a ribbon of 40-50mm. Clay contents between 30%-35% and with silt 25% or more.

Clayey sand: Slight coherence, sand grains of medium size, sticky when wet, many sand grains stick to fingers. Will form a minimal ribbon of 5-15 mm, discolours fingers with clay stain. Clay contents between 5%-10%.

Sandy clay loam: Strongly coherent bolus, sandy to touch, medium size grains visible in finer matrix, will form ribbon of 25 – 40mm. Clay contents between 20% - 30%.

Light clay: Plastic bolus, smooth to touch, slight resistance to shearing between thumb and fore finger, will form ribbon of 50 – 75 mm. Clay content between 35% - 40%.

Heavy Clay: Smooth plastic bolus, handles like stiff plasticine, can be moulded into rods without fracture, has firm resistance to ribboning shear. Will form a ribbon of 75 mm or more. Clay contents of 50% or more.

Sandy loam: Bolus coherent but very sandy to touch, will form a ribbon of 15-25 mm, dominant sand grains are of medium size and are readily visible. Clay contents between 10% - 20%.

Clay loam: Coherent plastic bolus, smooth to manipulate and will form a ribbon of 40 – 50 mm. Clay contents between 30% and 35%.

Light medium clay: Plastic bolus, smooth to touch, slight to moderate resistance to ribboning shear, will form a ribbon of about 75 mm.

Peat: Brownish or blackish fibrous substance produced by anaerobic decay of vegetation and found in boggy areas.

Comments: Any comments regarding soil structure, surface materials, rock contents or other soil attributes.

Source of water supply: The main sources of water that enter the water body, most commonly a combination of catchment run-off, groundwater and rainfall.

Rainfall estimate: Isohyet information gained from the Bureau of Meteorology for annual average rainfall. Also easily accessed by overlaying GIS isohyet information on the wetlands.

Dominant riparian vegetation association: Describes the dominant or co-dominant over storey species and understorey species.

Significance – why the wetland is important: Details any significance of the wetland within its regional setting or ecological function.

Noteworthy flora: Threatened species: threatened flora at national or State level that occur on the site (includes any threatened species identifies under national or State legislation, ANZECC lists or action plans). Composition information on the composition of any plant species or communities for which the wetland is particularly important.

Noteworthy fauna: Threatened species: list of threatened fauna at national or State level that are present at the site. (includes any threatened species identifies under national or State legislation, ANZECC lists or action plans). Composition: information regarding composition of important fauna that may inhabit the wetland permanently or seasonally including migratory species. An indication of population sizes, breeding colonies, migration stopovers is also given where available.

Land Degradation: disturbances or threats are defined as any direct or indirect human activities at the site or in the site or in the catchment area that may have a detrimental effect on the ecological character of the wetland. The effect may be a low level disturbance (eg low level grazing) or a major threat (eg water diversion schemes). Examples include disturbance by stock, water extraction, river regulation, siltation, salinity, urban development, drainage, pollution, excessive human activity, and impact of invasive species. There are several typical types of land degradation that may affect wetlands; these are listed as a standard template within the survey.

- Access tracks
- Cleared land
- Firebreaks
- Off-Road vehicles
- Rubbish dumping
- Pollution
- Pest vertebrate presence
- Altered water flows
- Borrow pits/Quarry
- Drains
- Fence lines
- Power lines
- Slashing/clearance
- Watering points
- Grazing damage
- Wetland drainage

Comments: Clarification of the severity of the threats and other comments.

Potential disturbances: Any threats that could impact on the wetland in the future, for example, a planned change in adjoining land use.

Conservation measures taken and measures proposed: Details of conservation measures being undertaken at the site (for example, fencing or revegetation) and the names of any protected areas established at or around the wetland. Details of management plans for the site and status of the site in terms of National Estate, Ramsar or World Heritage listing can also be included.

Current scientific research: Outline any research activities by scientific bodies such as CSIRO or universities or other recognised research organisations.

Current conservation education: Documents the existence of interpretive facilities, use of the area by school groups or other organisations.

Recreation and tourism: Records recreation facilities present on the site and if there are tourism uses or values with the site.

Social and Cultural Values: Documents pre-European and European historical and social values of the site.

References: Lists relevant references associated with the site.

Aquatic Vegetative Classes

This parameter provides a snap shot of aquatic vegetation structure, vegetation classes recorded include:

- Algal
- Aquatic moss
- Rooted vascular
- Floating leaved
- Floating vascular
- Unknown submergent
- Unknown surface

Aquatic Flora Sample

Records any aquatic vegetation sample collected with the wetland number, wetland name, GPS location, and habitat location. Samples should be placed into a snap lock plastic bag with water and labels placed inside and outside the bag. If field trips are longer than for one day, samples should be dried and pressed that day. Pressed samples will need the paper changed every other day to avoid the specimen rotting. The Plant Biodiversity Centre can assist with plant identification and a good reference books for aquatic plant identification include Sainty G.R (1994) *Waterplants in Australia* and Nick Romanowski (1998). *Aquatic and wetland plants*.

Aquatic Fauna Sample

Collecting Sample Material: This is achieved by one of two methods.

1. **Plankton Net.** This is a small round net of approx 200-250um pore size. Attached to the net is a length of string. One end of the string is tied to the plankton net. The other should be tied to the user of the net. The net is thrown into the water and is pulled through the water, towards the user. This procedure is done several times until sufficient sample material is collected.

If the water is too shallow then the net may be used by holding the metal rim and scooped through the water until sufficient sample material is collected in the plastic vial at the end of the net.

The sample material is then transferred from the sample vial to another vial, and 40% formalin is added to the sample to make up a mixture of approx.10% Formalin and 90% sample fluid.

2. **Hand Net.** The use of the hand net is a simple method. This is used in weedy habitats or shallow areas where the plankton net may become stuck or lost on objects. The net is used to prod at the base of vegetation and is run closely to the bottom of shallow water bodies. This is done over approx. 10-20 metres distance to obtain a representative sample from the water body.

The material in the net then is transferred to one or two sample vials.

Washing the Nets: This is done between sampling of each site in an attempt to clean the net of invertebrates to stop cross "contamination" of samples between sites. Cleaning of the net between sites is achieved by simply washing the net in the water of the site, without the sample vial attached. Ideally this should be done before and after a sample is taken.

Labelling Sample Vials: This is extremely important and should be done in a legible manner. The label should clearly and legibly be written upon and information recorded should be clearly stated.

Information recorded includes:

- Wetland number
- Name of wetland
- Date of Sample
- Grid Reference of site
- Method of collection (hand net or plankton net).

Material preservation of sample: material collected should be preserved with approximately 10% formalin as soon as possible after the sample is collected. this stops decomposition of the sample and also the invertebrates from eating each other. Formalin is used as microinvertebrates become soft and mushy, denaturing in the 70% alcohol solution used to preserve macroinvertebrates and vertebrates.

The formalin supplied is 40% by volume. This should be added to the sample material to make up approximately 10% formalin and 90% sample material. This is only a guide and should be used by the sampler with a degree of care.

Formalin is a dangerous chemical and is a fixative. Do not inhale it and wash your hands after use thoroughly. Read instructions on the label before use.

Water phys-chemistry

Five standard water chemistry parameters are collected; these are described below.

PH : is recorded using a Hanna HI 9025 pH meter. This meter is simply placed into the water body (ensuring that the probes do not touch the substrate) the reading is then given on the display. Calibration and maintenance of the pH unit is often needed, these procedures are outlined in the pH manual supplied with the unit.

Conductivity: is recorded using a Hanna HI 9635 meter. This meter can measure in the 0 to 199 $\mu\text{S}/\text{cm}$ range. It can be used to measure any sample from deionised water to highly saline water. The meter is placed in the water body ensuring that the probe does not touch the substrate and the reading is given on the display when stable. Calibration and maintenance procedures required are outlined in the manual.

Turbidity: is recorded using a Hanna HI 93703 portable microprocessor turbidity meter. The unit is designed to perform measurements according to the ISO 7027 International Standard. The instrument functions by passing a beam of light through a vial containing the sample being measured. A sensor, positioned at 90° with respect to the direction of light, detects the amount of light scattered by the undissolved particles present in the sample. These readings are given in NTU units. The manual accompanying the unit outlines measurement, calibration and maintenance procedures.

Dissolved O₂: is recorded using a Hanna HI 9142 dissolved oxygen meter. Dissolved oxygen is indicated in tenths of parts per million (ppm=mg/l). The dissolved oxygen probe has a membrane covering the polarographic sensors and a built in thermistor for temperature measurements and compensation. The thin permeable membrane isolates the sensor elements from the testing solution, but allows oxygen to enter. When a voltage is applied across the sensor, oxygen that has passed through the membrane reacts causing current to flow, allowing the determination of oxygen content.

Water Temperature: Water temperature is read from the pH or Dissolved Oxygen meter.

Rapid assessment (conservation values)

The rapid assessment component of the survey provides a quick snap shot of the vegetation associations and condition within different riparian habitats. Other parameters such as aquatic fauna and wetland condition is also recorded.

Aquatic Fauna: is a subjective score according to the abundance and diversity of invertebrates, fish and birds located within or on the water body.

- Low (1) refers to no fish or birds with little or no invertebrate presence.
- Moderate (3) refers to some bird presence and good vertebrate abundance, fish usually are not present.
- High (5) indicates good bird and invertebrate diversity and abundance with the presence of fish.

Aquatic Vegetation: records the abundance and diversity of aquatic vegetation.

- Low (1) indicates no or very little aquatic vegetation,
- Moderate (3) indicates some aquatic vegetation cover either in the form of floating or rooted vegetation,
- High (5) indicates good diversity of aquatic vegetation with a range of rooted vegetation such as reeds and rushes and floating vegetation such as water ribbons.

Riparian Vegetation

Riparian vegetation is assessed at three zones, these are the

- toe of bank (Low water – High water),
- bank (high water mark – buffer) and
- buffer (top of the bank to the buffer which can extend between 10 meters and 100 meters).

The vegetation association for each zone consists of the dominant or co-dominant over storey species and understorey species. The vegetation association for each zone is scored considering the level of disturbance and vegetation cover within each zone.

The vegetation association scoring system consists of:

- Degraded or no vegetation (1) with less than 30% vegetation cover, with high level of disturbance.
- Natural (3) between 30 – 75% vegetation cover, with little disturbance.
- Intact (5) over 75% vegetation cover with little or no disturbance.

Wetland Condition

The wetland score should reflect the previous rapid assessment scores for aquatic fauna, aquatic vegetation and riparian vegetation values. The combination of these values and the interpretation of others parameters recorded during the survey (such as land degradation

and water chemistry) form the basis of the wetland condition score.

The wetland score consists of:

- Degraded (1) those sites that have a high level of disturbance and received low rapid assessment scores.
- Natural (3) those sites that have little disturbance, received moderate to high rapid assessment scores and that are sites usually protected within the reserve system or by private conservation.
- Pristine (5) those sites with no obvious sign of disturbance, scored very highly in the rapid assessment and are formally conserved within the reserve system.

Photographic records

Photographic records for this survey have been recorded by digital camera in JPEG format. The following records should be kept when using a digital camera. It is recommended that a Laptop computer be taken on surveys for ease of cataloguing photographs.

All pictures need the following information recorded:

- Picture number and wetland number
- Direction and other features if relevant.

ANZECC wetland criteria

This criteria is used for determining nationally important wetlands in Australia, and form the bases for the inclusion in the Directory. These criteria are those agreed to by the ANZECC Wetlands Network in 1994.

A wetland may be considered nationally important if it meets at least one of the following criteria:

- It is a good example of a wetland type occurring within a biogeographic region in Australia.
- It is a wetland which plays an important ecological or hydrological role in the natural functioning of a major wetland system/complex.
- It is a wetland which is important as the habitat for animal taxa at a vulnerable stage in their life cycles, or provides a refuge when adverse conditions such as drought prevail.
- The wetland supports 1% or more of the national populations of any native plant or animal taxa.
- The wetland supports native plant or animal taxa or communities which are considered endangered or vulnerable at the national level.
- The wetland is of outstanding historical or cultural significance.

Many of the sites in the Directory meet more than one of the criteria. Application of the criteria to individual wetland sites involves a degree of subjectivity. Not only may certain aspects of a site's significance be interpreted differently by different investigators, but information gaps often exist which make it difficult to judge whether or not a site meets a particular criterion.

The Interim Biogeographic Regionalisation for Australia (IBRA) is used as the framework for applying Criterion 1, which identifies wetlands that are unique or representative within a biogeographic region in Australia.

Wetland type

The definition of a wetland used in the survey is one adopted by the Ramsar convention under Article 1.1.

“Wetlands are areas of marsh, fen, peatland or water, whether natural or artificial, permanent or temporary, with water that is static or flowing, fresh, brackish or salt, including areas of marine water the depth of which at low tide does not exceed six meters.”

Within this definition, the wetland classification system used in the Directory of Important Wetlands identifies 40 different wetland types in three categories: A-Marine and Coastal Zone wetlands, B-Inland wetlands, and C-Human-made wetlands. This wetland survey does not include Category A – Marine and Coastal Zone wetlands .

The system is based on that used by the Ramsar Convention in describing Wetlands of International Importance, but was modified slightly to suit the Australian situation.

A – Marine and Coastal Zone wetlands

- Marine waters - permanent shallow waters less than six metres deep at low tide, includes sea bays straits
- Subtidal aquatic beds, includes kelp beds, sea-grasses, tropical marine meadows
- Coral reefs
- Rocky marine shores, includes rocky offshore islands, sea cliffs
- Sand, shingle or pebble beaches, includes sand bars, spits, sandy islets
- Estuarine waters, permanent waters of estuaries and estuarine systems of deltas
- Intertidal mud, sand or salt flats
- Intertidal marshes, includes salt marshes, salt meadows, saltings, raised salt marshes, tidal brackish and freshwater marshes
- Intertidal forested wetlands, includes mangrove swamps, nipa swamps, tidal freshwater swamps forests
- Brackish to saline lagoons and marshes with one or more relatively narrow connections with the sea
- Freshwater lagoons and marshes in the coastal zone
- Non tidal freshwater forested wetlands

B – Inland wetlands

- Permanent rivers and streams includes waterfalls
- Seasonal and irregular rivers and streams
- Inland deltas(permanent)
- Riverine floodplains, includes river flats, flooded river basins, seasonally flooded grassland, savanna and palm savanna
- Permanent freshwater lakes (8 ha) includes large oxbow lakes
- Seasonal/intermittent freshwater lakes (>8 ha) floodplain lakes
- Permanent saline /brackish lakes
- Seasonal/intermittent saline lakes
- Permanent freshwater ponds (<8 ha) marshes and swamps on inorganic soils, with emergent vegetation waterlogged for at least most of the growing season
- Seasonal/intermittent freshwater ponds and marshes on inorganic soils includes sloughs, potholes, seasonally flooded meadows, sedge marshes
- Permanent saline/brackish marshes
- Seasonal saline marshes
- Shrub swamps, shrub dominated freshwater marsh, shrub carr, alder thicket

- on inorganic soil
- Freshwater swamp forest, seasonally flooded forest, wooded swamps, on inorganic soils
- Pearlands, forest, shrubs or open bogs
- Alpine and tundra wetlands: includes alpine meadows, tundra pools, temporary waters from snow melt
- Freshwater springs, oasis and rock pools
- Geothermal wetlands
- Inland, subterranean karst wetlands

C- Human-made wetlands

- Water storage areas; reservoirs, barrages, hydro-electric dams, impoundment's (generally over 8 ha).
- Ponds; includes farm ponds, stock ponds, small tanks; (generally below 8 ha).
- Aquaculture ponds; fish ponds shrimp ponds
- Salt exploitation, salt pans, salines
- Excavations; gravel pits; borrow pits, mining pools.
- Wastewater treatment areas; sewage farms, settling ponds, oxidation basins.
- Irrigated land; includes irrigation channels and rice fields, canals, ditches
- Seasonally flooded arable land, farm land
- Canals

Appendix 2 Invertebrate records for surveyed wetlands

Wetland reference number	Family	Genus	Species
S5070 - Breakneck River		<i>Cladotanytarsus</i>	sp
		<i>Paraterina</i>	
		<i>Tanytarsus</i>	sp
		<i>Culex</i>	sp
		<i>Austrochiltonia</i>	australis
		<i>Eucyclops</i>	cf baylyi
		<i>Eucyclops</i>	sp
		<i>Macrocylops</i>	sp
		<i>Paracyclops</i>	sp
		<i>Calamoecia</i>	ampulla
		<i>Gomphodella</i>	sp.
		<i>Cloeon</i>	sp
		<i>Notalina</i>	sp
		<i>Oxythira</i>	cf columba
	<i>Chironominus</i>	sp	
S5030 - Discovery Lagoon	Chironmidae	<i>Parachironomus</i>	sp
		<i>Mytilocypris</i>	tasmanica
S5066 - Six Mile Lagoon 2	Chironmidae	<i>Tanytarsus</i>	sp
	Chironmidae	<i>Procladius</i>	sp
	Corixidae	<i>Miconecta</i>	sp
	Notonectidae	<i>Enithares</i>	sp
	Lestidae	<i>Austrolestes</i>	sp
	Lestidae	<i>Austrolestes</i>	cingulatus
	Lestidae	<i>Austrolestes</i>	cingulatus
	Ceinidae	<i>Austrochiltonia</i>	australis
	Cyclopidae	<i>Australocyclops</i>	australis
	Cyclopidae	<i>Australocyclops</i>	australis
	Cyclopidae	<i>Diacyclops</i>	bisetosus
	Chironmidae	<i>Ablabesmyia</i>	sp
	Corixidae	<i>Miconecta</i>	sp
		<i>Triplectides</i>	australis
		<i>Tricoptera</i>	oecetis
	Notonectidae	<i>Enithares</i>	sp
	Cyclopidae	<i>Mesocyclops</i>	sp
	Notodromadidae	<i>Newnhamia</i>	fenestrata
		<i>Calamoecia</i>	ampulla
		<i>Boeckella</i>	symmetrica
	<i>Simocephalus</i>	sp	
	<i>Anisops</i>	sp	
	<i>Kennethia</i>	sp	
	<i>Candonocypris</i>	sp	
S5056 - Six Mile Lagoon	Cyclopidae	<i>Mesocyclops</i>	sp
	Cyclopidae	<i>Mixocyclops</i>	sp.
	Centropagidae	<i>Calamoecia</i>	ampulla
	Centropagidae	<i>Boeckella</i>	symmetrica
	Cyprididae	<i>Candonocypris</i>	sp
	Notodromadidae	<i>Kennethia</i>	sp
	Notodromadidae	<i>Newnhamia</i>	fenestrata
	Notodromadidae	UNIDENTIFIED	sp
	Daphniidae	<i>Daphnia</i>	cf carinata
	Daphniidae	<i>Simocephalus</i>	sp
	Chironmidae	(Pupae)	
	Corixidae	<i>Miconecta</i>	sp
	Corixidae	<i>Miconecta</i>	sp
	Centropagidae	<i>Calamoecia</i>	ampulla
	Centropagidae	<i>Boeckella</i>	symmetrica


Wetland reference number	Family	Genus	Species	
S5031 - Salt Lagoon	Ceratopogonidae			
	Ceratopogonidae			
	Cyclopidae	<i>Metacyclops</i>	<i>mortoni</i>	
	Cyprididae	<i>Diacypris</i>	<i>spinosa</i>	
	Notodromadidae	UNIDENTIFIED	sp	
	Branchipodidae	<i>Parartemia</i>	<i>zeitziana</i>	
	Daphniidae			
	Corixidae			
	Oniscidae	<i>Haloniscus</i>		
	S5018 - Rush Lagoon	Carabidae	(larvae)	
Dyticidae		<i>Necterosoma</i>	sp	
Hydrophilidae		<i>Berosus</i>	sp.	
Chironmidae		<i>Chironominus</i>	sp	
Chironmidae		<i>Paratanytarsus</i>	sp	
Chironmidae		<i>Procladius</i>	sp	
Corixidae		<i>Sigara</i>	sp	
Notonectidae		<i>Anisops</i>	sp	
Lestidae		<i>Austrolestes</i>	<i>annulosa</i>	
Lestidae		<i>Austrolestes</i>	<i>leda</i>	
Aeshnidae		<i>Aeshnia</i>	<i>brevistylia</i>	
Ceinidae		<i>Austrochiltonia</i>	<i>australis</i>	
Cyclopidae		<i>Australocyclops</i>	<i>australis</i>	
Cyclopidae		<i>Mixocyclops</i>	sp.	
Cyclopidae		<i>Mesocyclops</i>	sp	
Centropagidae		<i>Boeckella</i>	sp	
Limnocytheridae		<i>Mytilocypris</i>	<i>ambigua</i>	
Cyprididae		<i>Diacypris</i>	<i>spinosa</i>	
Notodromadidae		UNIDENTIFIED	sp	
Daphniidae		<i>Simocephalus</i>	sp	
Macrothricidae		<i>Macrothrix</i>	sp	
Chironmidae		<i>Tanytarsus</i>	sp	
Corixidae		<i>Sigara</i>	sp	
Notonectidae		<i>Anisops</i>	sp	
Cyclopidae		<i>Mesocyclops</i>	sp	
Cyclopidae		<i>Mesocyclops</i>	sp	
Cyclopidae		<i>Mesocyclops</i>	sp	
Cyclopidae		<i>Metacyclops</i>	<i>mortoni</i>	
		<i>Alboa</i>	<i>worooa</i>	
Limnocytheridae		<i>Mytilocypris</i>	<i>tasmanica</i>	
Cyprididae		<i>Diacypris</i>	<i>spinosa</i>	
Macrothricidae		<i>Macrothrix</i>	sp	
S5071 - Grassdale Lagoon		Chironmidae	<i>Chironominus</i>	sp
		Chironmidae	<i>Dicrotendipes</i>	sp
		Chironmidae	<i>Tanytarsus</i>	sp
		Chironmidae	<i>Procladius</i>	sp
		Chironmidae	(Pupae)	
		Culicidae	<i>Culex</i>	sp.
			<i>Enithares</i>	<i>woodwardi</i>
		Ceinidae	<i>Austrochiltonia</i>	<i>australis</i>
		Cyclopidae	<i>Mesocyclops</i>	sp
		Centropagidae	<i>Boeckella</i>	<i>symmetrica</i>
		Chydoridae	<i>Dunhevedia</i>	<i>crassa</i>
	Chydoridae	<i>Alona</i>	sp	
	Ilyocryptidae	<i>Ilyocryptus</i>	sp	
	Daphniidae	<i>Simocephalus</i>	<i>elizabethae</i>	
	Macrothricidae	<i>Macrothrix</i>	sp	
	Parastacidae			
	Simuliidae	<i>Cnephia</i>	sp	
	Centropagidae	<i>Calamoecia</i>	<i>australis</i>	
	Centropagidae	<i>Boeckella</i>	<i>symmetrica</i>	
	Chydoridae	<i>Dunhevedia</i>	<i>crassa</i>	
	Chydoridae	<i>Alona</i>	sp	
	Daphniidae	<i>Simocephalus</i>	<i>elizabethae</i>	
	Macrothricidae	<i>Macrothrix</i>	sp	
	Parastacidae			
		<i>Dicrotendipes</i>	sp	
		<i>Plea</i>	sp	
		<i>Hespercordulia</i>	sp	

Wetland reference number	Family	Genus	Species
S5064 - Duck Lagoon	Chironmidae	<i>Chironominus</i>	sp
	Chironmidae	<i>Dicrotendipes</i>	sp
	Chironmidae	<i>Tanytarsus</i>	sp
	Chironmidae	(Pupae)	
	Corixidae	<i>Agraptocorixa</i>	sp
	Corixidae	<i>Agraptocorixa</i>	sp
	Corixidae	<i>Miconecta</i>	sp
	Notonectidae	<i>Enithares</i>	<i>woodwardi</i>
	Cyclopidae	<i>Australocyclops</i>	<i>australis</i>
	Cyclopidae	<i>Australocyclops</i>	<i>australis</i>
	Cyclopidae	<i>Mesocyclops</i>	sp
	Notodromadidae	UNIDENTIFIED	sp
	Macrothricidae	<i>Macrothrix</i>	sp
	Hydrobiidae	<i>Fluvidona</i>	sp
	Hydrophilidae	<i>Berosus</i>	sp.
	Chironmidae	<i>Chironominus</i>	sp
	Chironmidae	<i>Dicrotendipes</i>	sp
	Chironmidae	<i>Procladius</i>	sp
	Ceratopogonidae		
	Notonectidae	<i>Anisops</i>	sp
	Notonectidae	<i>Enithares</i>	sp
	Aeshnidae	<i>Hemianax</i>	<i>papuensis</i>
	Oxygastridae	<i>Hesperocordula</i>	sp
	Ceinidae	<i>Austrochiltonia</i>	<i>australis</i>
	Cyclopidae	<i>Mesocyclops</i>	sp
	Centropagidae	<i>Boeckella</i>	<i>symmetrica</i>
	Centropagidae	<i>Boeckella</i>	sp
	Notodromadidae	UNIDENTIFIED	sp
	Ilyocryptidae	<i>Ilyocryptus</i>	sp
		<i>Tricoptera</i>	<i>oecetis</i>
	Centropagidae	<i>Calamoecia</i>	<i>clittelata</i>
	Centropagidae	<i>Boeckella</i>	<i>symmetrica</i>
S5065 - Cygnet River Lagoon	Dyticidae	<i>Chostonectes</i>	sp
	Chironmidae	<i>Chironominus</i>	sp
	Corixidae	<i>Agraptocorixa</i>	sp
	Corixidae	<i>Miconecta</i>	sp
	Notonectidae	<i>Anisops</i>	sp
	Ceinidae	<i>Austrochiltonia</i>	<i>australis</i>
	Cyclopidae	<i>Australocyclops</i>	<i>similis</i>
	Cyclopidae	<i>Mesocyclops</i>	sp.
	Centropagidae	<i>Boeckella</i>	<i>symmetrica</i>
	Notodromadidae	UNIDENTIFIED	sp
	Daphniidae	<i>Ceriodaphnia</i>	sp
	Daphniidae	<i>Daphnia</i>	sp
	Daphniidae	<i>Simocephalus</i>	<i>cf expinosus</i>
Macrothricidae	<i>Macrothrix</i>	sp	
S5020 - Big White Salt Lagoon	Branchipodidae	<i>Parartemia</i>	<i>zeitziana</i>
S5013 - Greenfields Wetland	Cypridae	<i>Diacypris</i>	<i>spinosa</i>
	Branchipodidae	<i>Parartemia</i>	<i>zeitziana</i>
S5043 - Emanuel Swamp		<i>Keratella</i>	sp.
	Ceinidae	<i>Austrochiltonia</i>	<i>australis</i>
	Cyclopidae	<i>Apocyclops</i>	<i>annulosus</i>
	Cyclopidae	<i>Metacyclops</i>	<i>mortoni</i>
	Centropagidae	<i>Calamoecia</i>	<i>clittelata</i>
	Centropagidae	<i>Boeckella</i>	<i>triarticulata</i>
Limnocytheridae	<i>Mytilocypris</i>	<i>tasmanica</i>	
S5000 - Lashmar Lagoon	Cyclopidae	<i>Mesocyclops</i>	sp
	Cypridae	<i>Diacypris</i>	<i>spinosa</i>
	Notodromadidae	UNIDENTIFIED	spp
	Macrothricidae	<i>Macrothrix</i>	sp
S5035 - Salt Lake	Centropagidae	<i>Calamoecia</i>	<i>salina</i>
	Daphniidae	<i>Daphniopsis</i>	sp

Wetland reference number	Family	Genus	Species
S5021 - White Lagoon1	Cyclopidae	<i>Metacyclops</i>	<i>mortoni</i>
	Centropagidae	<i>Calamoecia</i>	<i>salina</i>
	Limnocytheridae	<i>Mytilocypris</i>	<i>ambiguosa</i>
	Notodromadidae	UNIDENTIFIED	Spp
	Branchipodidae	<i>Parartemia</i>	<i>zeitziana</i>
S5062 - Nepean Bay Lagoon	Centropagidae	<i>Calamoecia</i>	<i>salina</i>
	Limnocytheridae	<i>Limnocythere</i>	<i>porhyretica</i>
	Cyprididae	<i>Diacypris</i>	<i>spinosa</i>
S5006 - Milky's Lake	Cyclopidae	<i>Metacyclops</i>	<i>mortoni</i>
	Centropagidae	<i>Calamoecia</i>	<i>salina</i>
	Cyprididae	<i>Diacypris</i>	<i>spinosa</i>
	Notodromadidae	UNIDENTIFIED	Spp
	Branchipodidae	<i>Parartemia</i>	<i>zeitziana</i>
	Oniscidae	<i>Haloniscus</i>	<i>searli</i>
S5044 - Destree Swamp	Notonectidae	<i>Anisops</i>	sp
	Ceinidae	<i>Austrochiltonia</i>	<i>australis</i>
	Limnocytheridae	<i>Mytilocypris</i>	<i>tasmanica</i>
	Cyprididae	<i>Diacypris</i>	<i>spinosa</i>
S5042 - Wangara Lagoon	Ceratopogonidae		
	Cyprididae	<i>Diacypris</i>	<i>spinosa</i>
	Notodromadidae	UNIDENTIFIED	Spp
	Branchipodidae	<i>Parartemia</i>	<i>zeitziana</i>
S5063 - Salt Lake 2	Ceratopogonidae		
	Ceratopogonidae		
	Notodromadidae	<i>Alboa</i> UNIDENTIFIED	<i>worooa</i> spp
S5053 - Murrays Lagoon	Notonectidae	<i>Anisops</i>	sp
	Ceinidae	<i>Austrochiltonia</i>	<i>australis</i>
	Cyclopidae	<i>Neocyclops</i>	<i>petkovis</i>
		<i>Alboa</i>	<i>worooa</i>
	Limnocytheridae	<i>Mytilocypris</i>	<i>tasmanica</i>
	Cyprididae	<i>Diacypris</i>	<i>spinosa</i>
S5050 - White Lagoon	Stratiomyidae	<i>Odontomyia</i>	sp
	Notonectidae	<i>Enithares</i>	sp
	Lestidae	<i>Austrolestes</i>	sp
	Cyclopidae	<i>Australocyclops</i>	<i>similis</i>
	Cyclopidae	<i>Australocyclops</i>	sp
	Centropagidae	<i>Calamoecia</i>	<i>ampulla</i>
		<i>Alboa</i>	<i>worooa</i>
		<i>Cypricercus</i>	sp
	Limnocytheridae	<i>Mytilocypris</i>	<i>ambiguosa</i>
	Cyprididae	<i>Diacypris</i>	<i>spinosa</i>
	Daphniidae	<i>Daphniopsis</i>	sp
Macrothricidae	<i>Macrothrix</i>	sp	
S5041 - Halls Road Swamp	Chironmidae	<i>Procladius</i>	sp
	Lestidae	<i>Austrolestes</i>	<i>annulosa</i>
	Ceinidae	<i>Austrochiltonia</i>	<i>australis</i>
	Cyclopidae	<i>Apocyclops</i>	<i>annulosus</i>
	Centropagidae	<i>Calamoecia</i>	<i>clittelata</i>
	Centropagidae	<i>Boeckella</i>	<i>triarticulata</i>
		<i>Alboa</i>	<i>worooa</i>
	Limnocytheridae	<i>Mytilocypris</i>	<i>ambiguosa</i>
	Cyprididae	<i>Diacypris</i>	<i>spinosa</i>

Wetland reference number	Family	Genus	Species
S5055 - Birchmore Lagoon		<i>Alboa</i>	<i>worooa</i>
		<i>Mytilocypris</i>	<i>tasmanica</i>
		<i>Diacypris</i>	<i>spinosa</i>
		<i>Anisops</i>	<i>sp</i>
		<i>Macrothrix</i>	<i>sp</i>
		<i>Tanythrusus</i>	<i>sp</i>
		<i>Metacyclops</i>	<i>mortoni</i>
		<i>Mesocyclops</i>	<i>sp</i>







Map 1 Kangaroo Island Natural Heritage Trust Boundary



	Aboriginal Lands		Marks		Parliamentary
	Byron		Miscellaneous Properties		State Land
	Parliamentary Land		Murray Darling Basin		
			South-eastern Agricultural District		




Produced by: Resource Services
 Date: 2013
 Data source: 1:50,000 Scale and 1:250,000 Scale
 South Australian Department of Environment and Heritage
 Department for Environment and Heritage
 Project No: 0000000000
 Date: 2013
 Drawn by: [Name]
 Checked by: [Name]

Map 2 Surveied wetlands on Kangaroo Island




Produced by Russel Semman
 Data source In Motion and Data Analysis Branch, Planning SA, Department of Transport, Urban planning and the Arts.
 Department for Environment and Heritage
 Projection Decimal degrees
 Datum GDA 14
 Compiled February 2011

Map 3 Environmental Associations on Kangaroo Island




Produced by Russell Stearn
 Data source: Information and Data Analysis 30
 Bands, Planning SA, Department
 for Transport, Urban planning and
 the Arts.
 Department for Environment and
 Heritage
 Project on: District design
 Date: GDA 94
 Compiled: February 2002

Map 4 Key Biodiversity Areas on Kangaroo Island




Key Biodiversity Areas

	Fragmented Habitat Area
	Large Remnant Area
	Threatened Habitat Area





Produced by: Russell Stewart
 Date issued: July 2005 and 2010 updates
 Based on: Planning IIS, Stage 1 and 2
 for Strategic Planning and Management
 of IIS
 Department for Environment and
 Heritage
 Population: General Population
 Series: 2005-06
 Copyright: © 2005-06

Map 5 Geological features underlying surveyed wetlands on Kangaroo Island



Geology 1		Geology 2	
	Undifferentiated Carboniferous-Permian rocks		Bridge water Formation
	Undifferentiated alluvial/fluvial sediments		Kangaroo Group
	Undifferentiated lacustrine/ Playa sediments		



Produced by: Russell Steiner
 Data source: Information and Data Analysis 30
 Branch, Planning SA, Department
 for Transport, Urban planning and
 the Arts.
 Department for Environment and
 Heritage
 Project on: Datasat digital
 Date: GDA 94
 Compiled: February 2002